

# MINIREVIEW

## Seasonal Changes in Adiposity: the Roles of the Photoperiod, Melatonin and Other Hormones, and Sympathetic Nervous System

TIMOTHY J. BARTNESS,<sup>1,\*†</sup> GREGORY E. DEMAS,\* AND C. KAY SONG\*

*Departments of \*Biology and of †Psychology, Neurobiology and Behavior Program, and Center for Behavioral Neuroscience, Georgia State University, Atlanta, Georgia 30303*

It appears advantageous for many non-human animals to store energy body fat extensively and efficiently because their food supply is more labile and less abundant than in their human counterparts. The level of adiposity in many of these species often shows predictable increases and decreases with changes in the season. These cyclic changes in seasonal adiposity in some species are triggered by changes in the photoperiod that are faithfully transduced into a biochemical signal through the nightly secretion of melatonin (MEL) via the pineal gland. Here, we focus primarily on the findings from the most commonly studied species showing seasonal changes in adiposity—Siberian and Syrian hamsters. The data to date are not compelling for a direct effect of MEL on white adipose tissue (WAT) and brown adipose tissue (BAT) despite some recent data to the contrary. Thus far, none of the possible hormonal intermediaries for the effects of MEL on seasonal adiposity appear likely as a mechanism by which MEL affects the photoperiodic control of body fat levels indirectly. We also provide evidence pointing toward the sympathetic nervous system as a likely mediator of the effects of MEL on short day-induced body fat decreases in Siberian hamsters through increases in sympathetic drive on WAT and BAT. We speculate that decreases in the SNS drive to these tissues may underlie the photoperiod-induced seasonal increases in body fat of species such as Syrian hamsters. Clearly, we need to deepen our understanding of seasonal adiposity, although, to our knowledge, this is the only form of environmentally induced changes in body fat where the key elements of its external trigger have been identified and can be traced to and through their transduction into a physiological stimulus that ultimately affects identified responses of white

adipocyte physiology and cellularity. Finally, the comparative physiological approach to the study of seasonal adiposity seems likely to continue to yield significant insights into the mechanisms underlying this phenomenon and for understanding obesity and its reversal in general.

[Exp Biol Med Vol. 227(6):363–376, 2002]

**Key words:** white fat; adipose tissue; brown fat; brown adipose tissue; thyroid; testosterone; estradiol; leptin; insulin; glucagon; glucocorticoid; prolactin; thermogenesis; reproduction; pineal

Obesity in humans is a disease of figuratively and literally enormous proportions. The secondary health consequences of obesity include cardiovascular disease, stroke, noninsulin-dependent diabetes mellitus, and carcinomas (reviewed in Ref. 1). Because of these secondary health consequences, there seems to be little survival advantage for humans to eat and store as much energy as possible in lipid reserves, especially when the food supply is relatively constant and plentiful as it is in most economically developed countries. It has been hypothesized that a contributing factor to the prevalence of obesity in contemporary society is the persistence of metabolically efficient systems for storage of energy as body fat (2). In contrast to humans, it appears advantageous for many non-human animals to store energy body fat extensively and efficiently because their food supply is more labile and less abundant than in their human counterparts. The level of adiposity in many of these species does not continuously increase across their life span, however; instead, body fat levels often shows predictable increases and decreases with changes in the season. Species exhibiting seasonal adiposity can be divided into two major categories. The first category includes species that respond to changes in the photoperiod (daylength; e.g., hamsters and voles), some of which will be

---

This work was supported in part by the National Institutes of Health (grant RO1 DK35254 and NRSA NS-10596 to G.E.D.) and by the National Institute of Mental Health (Research Scientist Development Award KO2 MH00841 to T.J.B.).

<sup>†</sup>To whom requests for reprints should be addressed at Department of Biology, Georgia State University, 24 Peachtree Center Avenue, NE, Atlanta, GA 30303. E-mail: bartness@gsu.edu

---

1535-3702/02/2276-0363\$15.00

Copyright © 2002 by the Society for Experimental Biology and Medicine

the primary focus of this review, whereas the second category includes species that respond to the timing of an endogenous “clock” of unknown location (e.g., ground squirrels, wood chucks, and marmots), that will not be our focus (reviewed in Ref. 3). Species in both categories appear to shift from the obese to the lean state effortlessly, unlike humans. We will focus on the possible mechanisms underlying photoperiod-induced seasonal adiposity cycles.

### **Animals Display Seasonal Changes in Physiology and Behavior, Especially in Temperate Zones**

Animals living in temperate zones frequently show seasonal changes in a host of physiological and behavioral responses. For example, the timing of reproduction is critical to many small rodent species living in temperate zones. Most of these species are reproductively quiescent in the late fall/winter, when ambient temperatures decline and food availability is typically low, and they resume reproductive activities in the early spring/summer when ambient temperature and the food supply are more favorable (reviewed in Ref. 4). Although photoperiod-induced changes in seasonal adiposity occur in a variety of mammalian species, they have been studied most extensively in Syrian (*Mesocricetus auratus*) and Siberian hamsters (*Phodopus sungorus*; reviewed in Refs. 5 and 6); therefore, these species will be highlighted throughout this review. The seasonally adaptive changes in lipid stored in white adipose tissue (WAT, the major energy storage depot for mammals; 7), and in thermogenesis by brown adipose tissue (BAT, an important site for heat generation in mammals; 8), often takes weeks or months to be manifested fully. Therefore, it appears that natural selection has favored animals that accurately begin the appropriate changes in these tissues, as well as those more directly associated with reproduction, in anticipation of the forthcoming season.

The photoperiod is the critical environmental cue that triggers many seasonal responses and it is transduced into a biochemical signal by the pineal gland. The change in the photoperiod (daylength) is the most noise-free signal in the environment indicating the forthcoming season (9). Therefore, by simply changing the photoperiod in the laboratory, the entire constellation of photoperiodic responses in nature can be conveniently mimicked and manipulated (4, 10). For example, hamster and vole species will show the full complement of short-day (SD) “winter-like” responses if the daylength is significantly shortened from a long-day (LD) “summer-like” photoperiod (11–15). Procedurally, this often is done by switching animals from a 16:8-hr or 14:10-hr light:dark cycle to an 8:16-hr or 10:14-hr light:dark cycle.

This photoperiod cue is received by the retinal ganglion cells and is transmitted through a multisynaptic pathway to the pineal gland (16). Interruption of this circuit at any point, including the eyes or the pineal, blocks the effects of SDs on the reproductive system (17) and most other sea-

sonal responses, including the seasonal changes in adiposity (11, 18). For example, pinealectomy blocks the photoperiod-dependent changes in body and lipid mass in Syrian hamsters (11), although evidence for a photoperiod-induced, pineal-independent effect also has been reported (11, 19). Pinealectomy also blocks the seasonal changes in body and lipid mass of Siberian hamsters (18, 20, 21) and meadow voles (22, 23). Once the neurochemical interpretation of the photoperiod reaches the pinealocytes via its sympathetic nervous system (SNS) innervation, it is transduced into an endocrine signal in the form of the rhythmic secretion of the melatonin (MEL) into the blood. Pineal and plasma MEL concentrations are at their nadir during the light phase of the photocycle and at their peak during the dark phase (reviewed in Refs. 24 and 25). This pattern of MEL synthesis and secretion results from the regulation of the pineal gland by an endogenous circadian oscillator that is entrained to the light:dark cycle (26). Thus, the duration of the night is faithfully coded into the duration of MEL secretion and serves to trigger seasonal responses—long MEL secretion durations signal “fall/winter,” whereas short MEL secretion durations signal “spring/summer” (reviewed in Ref. 25).

The peak duration of MEL secretion is its critical feature for triggering SD responses. This was initially shown by giving daily subcutaneous injections of MEL to LD-housed, pineal-intact Syrian or Siberian hamsters ~2 to 3 hr before lights out (i.e., the “timed afternoon injection paradigm”; 27). Hamsters treated in this manner exhibited SD-like seasonal changes in body and lipid mass (11, 28, 29) and in reproductive status (30–32). Why does this happen? One notion is that the exogenously administered MEL summates with the nocturnally secreted endogenous MEL to lengthen the peak duration of circulating MEL, thereby approximating those occurring naturally in SDs (reviewed in Ref. 25). This critical nature of the duration of MEL secretion for triggering photoperiod-induced changes in adiposity and reproductive status was later unequivocally shown by giving LD-housed, pinealectomized Siberian or Syrian hamsters, or sheep, precisely timed daily subcutaneous MEL infusions instead of bolus injections (i.e., termed the “timed infusion paradigm”) so as to mimic the naturally occurring peak durations of circulating MEL in SDs (reviewed in Ref. 25). For example, long-duration MEL infusions triggered SD-like decreases in body mass and fat, and gonadal regression in adult male Siberian hamsters (21, 33, 34).

### **MEL Receptors Occur in the Brain and Periphery, but Stimulation of Specific Central Sites Appears Necessary for Seasonal Changes in Adiposity and Reproductive Status**

MEL brain receptors were first suggested by the binding of a radioactive MEL analogue (2-[<sup>125</sup>I]iodomelatonin; IMEL) to neural tissue (reviewed in Refs. 35–37). Only a relatively few brain MEL binding sites exist (38, 39), although MEL binds extensively to many peripheral tissues

(e.g., 40–44). IMEL binds in the brains of a wide range of mammals (e.g., sheep, deer, goats, rabbits, laboratory rats, Syrian, Siberian, and European hamsters, white-footed mice, golden-mantled ground squirrels, Guinea pigs, nine-banded armadillos, and humans), as well as non-mammals (Atlantic hagfish, lamprey, hedgehog skates, rainbow trout, clawed toads; reviewed in Ref. 35). Across many mammalian species, one brain and one peripheral site consistently show IMEL binding—the suprachiasmatic nucleus of the hypothalamus (SCN) and the pars tuberalis (PT) of the pituitary gland (45–48), respectively. Many of these species show binding in both the SCN and the PT, although there are a few seasonally breeding species where IMEL binding only is found in the PT (49, 50). Gene expression for one of the MEL receptor subtypes, the MEL  $1_a$  receptor (also known as the  $mt_1$  receptor) is present in the SCN and PT of Siberian hamsters and laboratory rats (51). The MEL  $1_a$  receptors of the SCN are important for the seasonal changes in reproductive status and adiposity in at least Siberian hamsters. Thus, daily SD-like MEL signals given via the timed-infusion paradigm to pinealectomized Siberian hamsters bearing SCN lesions (33, 34, 52), do not trigger SD-like decreases in body and WAT pad masses, as well as gonadal regression. In Syrian hamsters, however, SCN lesions do not block SD-like MEL-induced gonadal regression (53). In addition, infusions of MEL directly into the SCN of juvenile Siberian hamsters inhibit maturation of the reproductive system, mimicking the effects of SD exposure in these animals (54).

The SCN output pathways (efferents) important for relaying the photoperiod-encoded MEL signals to other brain/peripheral targets involved in seasonal responses are unclear at present (reviewed in Ref. 55). It appears MEL works via an intermediary, at least for the seasonal changes in adiposity, because WAT lipolysis (fat mobilization) and lipogenesis (fat synthesis) are not triggered in isolated adipocytes using physiological doses of MEL in Syrian hamsters, laboratory rats, and rabbits (56). According to a recent report, however, lipolysis is inhibited by MEL *in vitro* in laboratory rat inguinal, but not epididymal WAT (57). Furthermore, MEL can act directly on Siberian hamster BAT cells through a MEL-specific binding site (but not the MEL $_{1a}$  receptor isoform; 44), as well as modulating BAT mitochondrial cytochrome b gene expression directly (58). Thus, although there are some direct effects of MEL on its peripheral target adipose tissues via its receptors, the preponderance of evidence suggests that the MEL signal is received by brain MEL receptors.

The means by which the season-encoded MEL signal reaches peripheral tissues after stimulating central MEL receptors is not well understood. How stimulation of these receptors ultimately affects peripheral tissues is not certain at this time. One possible means by which MEL ultimately affects adipose tissues, gonads, hair follicles, and other peripheral endpoints that change seasonally involves projections from neurons within the SCN and/or other brain sites

that possess MEL $_{1a}$  receptors to these target tissues via a humeral intermediary. Alternatively, stimulation of the brain MEL receptors may trigger the seasonal changes in these tissues strictly through neural communication of the photoperiod-encoded MEL signals.

### Do the Hormones that Fluctuate Seasonally Mediate Photoperiod-Induced Changes in Adiposity?

Although there are many candidates for the hormonal intermediary controlling the photoperiod (MEL) effects on seasonal adiposity (for a more complete review, see Ref. 6), the supporting data for such a humoral intermediary are few. Here we discuss possible roles of gonadal steroids, prolactin, thyroid hormones, glucocorticoids, insulin, glucagon, and leptin in mediating seasonal adiposity in Siberian and Syrian hamsters.

**Gonadal Steroids.** Hamsters (12, 59–61), voles (15, 62), collared lemmings (63), and deer mice (64, 65) all regress their gonads in response to SD exposure; however, their body and lipid mass responses differ (see below). That is, SD-exposed collared lemmings (66), prairie voles (14), and Syrian hamsters (11, 67, 68), increase, whereas SD-exposed Siberian (12, 13, 61, 69, 70) and European hamsters (60), meadow voles (15, 71), and deer mice (72, 73) decrease their body and lipid mass. Across these and many other rodent species, the direction of the SD-induced change in adiposity can be predicted by the effects of gonadectomy on the body fat levels of LD-housed individuals of these species. This lead to the speculation that the effects of gonadectomy and the SD-induced “functional gonadectomy” on body fat are both primarily due to the decreases in testosterone and estrogen. For example, LD-housed female Syrian hamsters (74, 75), female collared lemmings (66), and male prairie voles (14) increase their body and lipid masses when they are gonadectomized as well as when they are “photoperiodically gonadectomized” by SDs (11, 14, 66). In contrast, Siberian hamsters (13, 76), male deer mice (77), meadow voles (71, 78), and European hamsters (60, 79) decrease their body and lipid masses when gonadectomized and also decrease body and lipid masses when exposed to SDs (60, 69, 79).

Because gonadectomized animals exposed to SDs have body fat increases or decreases that are similar to their SD-exposed gonad-intact counterparts, this obvious explanation for the effects of SD exposure on seasonal adiposity is not satisfactory. That is, the gonadectomy-induced decrease in body fat of LD-housed Siberian hamsters is decreased further when they are subsequently exposed to SDs (13), and the gonadectomy-induced increase in body fat of LD-housed Syrian hamsters is increased further when they are subsequently exposed to SDs (11). In addition, body and lipid mass (along with all other photoperiodic responses) revert to their characteristic LD values after transfer from SDs to LDs or after extended exposure to SDs (i.e., “spontaneous recrudescence”), and these changes in body and

lipid mass also are partly independent of the gonadal responses. Thus, when ovariectomized Syrian hamsters (28), castrated male European hamsters (60), castrated male meadow voles (71), and cold-exposed castrated male Siberian hamsters (81) are exposed to persisting SD exposure, their species-specific SD-induced increases or decreases in body and lipid mass are reversed despite being gonadectomized. In addition, if SD-housed castrated male Siberian hamsters at their body mass nadir are abruptly shifted to LDs, they show a nearly identical increase in body mass compared with SD-housed testes-intact hamsters experiencing identical changes in the photoperiod (82). Therefore, the effects of the photoperiod on gonadal steroid secretion and action are not primarily responsible for seasonal changes in body mass and fat in these animals.

**Prolactin.** All species that show photoperiod control of reproductive cycles, regardless of whether they are LD or SD breeders (83), show SD-induced decreased prolactin (PRL) serum concentrations (84). An important role for PRL in the photoperiodic regulation of lipid deposition or mobilization has not been clearly demonstrated, but PRL is capable of altering WAT lipid metabolism both *in vivo* and *in vitro* in laboratory rats (85, 86). Syrian and Siberian hamsters show the typical SD-induced decreases in serum PRL concentrations (87). Tests of the role of PRL in seasonal obesity have largely been negative. For example, injections of the dopamine receptor agonist bromocriptine (CB-154), an inhibitor of PRL secretion (88), designed to produce SD PRL levels in LD-housed Syrian or Siberian hamsters does trigger their species-specific SD increases or decreases in changes in lipid mass or WAT metabolism, respectively but not to the same degree as SDs (89). Conversely, LD pituitary explants or infusions of ovine PRL designed to produce LD serum concentrations of PRL in SD-housed Syrian and Siberian hamsters, respectively, do not revert them to their species-specific LD levels of body fat (89, 90). Therefore, at least in Syrian and Siberian hamsters, it does not appear that PRL per se is a significant mediator of photoperiod-induced changes in body and lipid mass. It should be noted, however, that a critical role for PRL in the development of photoperiod-induced seasonal changes in body fat (91–95) has been reported by one laboratory, although it is not supported by others (89, 96).

**Thyroid Hormones.** The pineal gland and MEL modulate the neuroendocrine-thyroid gland axis in several species and consequently, the secretion of thyroxine ( $T_4$ ) and triiodothyronine ( $T_3$ ) (reviewed in Ref. 97). SD exposure significantly decreases serum concentrations of  $T_4$  and  $T_3$  in Syrian hamsters (98, 99) and collared lemmings (100, 101) and more modestly in Siberian hamsters (102–104). These SD-induced reductions in thyroid hormones are somewhat surprising because “winter-like” SDs would typically be associated with decreases in ambient temperature in the wild, and most rodent species increase their secretion of thyroid hormones in response to cold exposure (105, 106). It may be that the neuroendocrine-thyroid axis is only

stimulated when animals actually experience the cold, but not in preparation for cold exposure. Indeed, SD exposure alone only produces a transient increase in  $T_3$  in Syrian hamsters with no enduring effects (107). The photoperiod via MEL does seem important in BAT thermogenesis, however, in that MEL treatment stimulates type-II thyroxine 5'-deiodinase in BAT, the enzyme responsible for local tissue conversion of  $T_4$  to  $T_3$  (105). Moreover, increases in deiodination of  $T_4$  facilitate BAT UCP-1 gene expression in the cold (106). Thus, thyroid hormones may be important for the SD increase in BAT thermogenesis, but seem to require the addition of cold to be maximally effective. The role of thyroid hormones in mediating the photoperiod-induced changes in adiposity, however, is less clear. For example, SD-housed male Siberian hamsters given daily  $T_4$  injections have LD-like, high serum  $T_4$  and  $T_3$  concentrations, yet show decreases in body, fat pad, and paired testes masses that are similar to those of SD-housed vehicle- and noninjected hamsters (104). Therefore, photoperiod/MEL-induced changes in thyroid hormones do not seem have a major role in the seasonal control of body and lipid mass, at least in these hamster species.

**Glucocorticoids.** The role of the glucocorticoids in the photoperiod/MEL-controlled alterations in energy balance has not been thoroughly investigated; however, the role of glucocorticoids in all aspects of energy balance, including food intake, lipid mobilization and storage, and thermogenesis is well established in several categories of rodent obesity (dietary, genetic, endocrine, and hypothalamic; 108). A relation between glucocorticoid secretion and the photoperiod has been shown for some of the species discussed above. For example, Syrian (99, 109) and Siberian hamsters (110), prairie voles (111), and collared lemmings (101) show seasonal changes in serum cortisol and/or corticosterone concentrations or changes in both (note that unlike laboratory rats [112, 113] or humans [114], hamsters have significant serum concentrations of both cortisol and corticosterone concentrations [99, 109, 110, 115]). The effects of adrenal glucocorticoids on WAT mass and function may be mediated by the Type II glucocorticoid receptor found in the cytosol of WAT adipocytes (116). Decreases in glucocorticoid secretion and/or decreases in activation of Type II glucocorticoid receptors via adrenalectomy is not feasible in Siberian hamsters because the additional loss of mineralocorticoids by this surgery does not trigger increases in salt appetite as it does in laboratory rats and mice, thereby leading to their death (T. Bartness and B. Goldman, unpublished data). Therefore, to study the possible role of glucocorticoids in hamsters a different tact is needed.

First, although glucocorticoids promote adiposity in the typically studied rodent obesity models, it seemed possible that they might produce an opposite effect in Siberian hamsters, given the obesity promoting effects of estrogen and testosterone in this hamster species (76, 117). Therefore, Type II glucocorticoid receptor blockade by RU-486 (RU-

486 is not an progesterin receptor antagonist in hamsters; 118) was used to produce a SD-like inhibition of glucocorticoid function in LD-housed hamsters. LD-housed Siberian hamsters treated chronically with RU-486 for several weeks did not decrease body and lipid mass as occurs when they are transferred from LDs to SDs (110). Thus, unlike the inhibition of obesity by Type II glucocorticoid receptor blockade using RU-486 in genetic- and diet-induced obese laboratory rats (119, 120), these receptors, and perhaps glucocorticoids in general, do not seem to be a primary factor in the photoperiodic control of seasonal body and lipid mass cycles in this species and perhaps in other photoperiodic species.

**Insulin and Glucagon.** Of the species showing photoperiod-induced change in adiposity, seasonal fluctuations of serum insulin concentrations only have been measured in Syrian and Siberian hamsters to our knowledge. Specifically, SD exposure decreases serum insulin concentrations in Siberian hamsters (115), but increases serum insulin concentrations in Syrian hamsters (92–94, 121). In Siberian hamsters, serum insulin concentrations decrease by 4-fold in SDs compared with LDs (115), consistent with the positive correlation between circulating insulin concentrations and level of adiposity in most mammals (reviewed in Ref. 122). Despite this relation, normal SD-induced decreases in body and lipid mass, food intake, and gonadal regression are seen in Siberian hamsters treated with streptozotocin to induce diabetes mellitus and then one of several levels of insulin replacement to produce a wide range of serum insulin concentrations (123). These data suggest that insulin status does not play a major role in the photoperiodic control of seasonal adiposity, at least for this species.

Insulin also has been manipulated experimentally in Syrian hamsters, but indirectly, to test for its role in SD-induced body fat increases (121). That is, the neural control of insulin secretion by the parasympathetic nervous system was eliminated by total subdiaphragmatic vagotomy. This treatment blocked the SD-induced increases in body and lipid mass characteristic of this species, but did not block gonadal regression (121). Interpretation of these data is difficult, however, because total subdiaphragmatic vagotomy causes a multitude of effects (124) in addition to the blockade of neurally released insulin. Therefore, the inability of vagotomized Syrian hamsters to increase body mass when exposed to SDs does not necessarily lend support for a role of insulin in the SD-induced obesity of this species.

Although the pancreatic and intestinal hormone glucagon is well known for its stimulation of WAT lipolysis and of BAT thermogenesis, to our knowledge, serum concentrations of glucagon have not been measured in any of the rodent species showing photoperiod-mediated changes in seasonal adiposity. It seems quite likely, however, that glucagon may play a significant role in the photoperiodic control of energy balance because it stimulates BAT thermogenesis (125), is an important hormone in the thermogenic response to the cold (126), and has receptors on WAT adi-

pocytes (127). Thus, in species showing SD-induced decreases in adiposity, such as Siberian hamsters, SD increases in glucagon secretion, which show a reciprocal relation with insulin (128), could increase lipid mobilization from WAT and also could increase the thermogenic capacity of BAT, whereas in species showing SD-induced increases in adiposity, such as Syrian hamsters, the opposite might occur. This intriguing notion remains to be tested.

**Leptin.** In virtually all mammalian species undergoing seasonal fluctuations in body fat, there are concomitant fluctuations in circulating leptin, a peptide hormone produced primarily, but not exclusively, by WAT (reviewed in Refs. 129 and 130). Because both leptin and its receptors are integral components of a hypothesized feedback system that regulates body fat levels (reviewed in Ref. 131), leptin also may be involved in seasonal control of body fat. According to the commonly accepted, leptin feedback system hypothesis, as an animal gets fatter, more leptin is secreted from the growing adipose depots. The resulting increases in concentrations of circulating leptin stimulate brain (and other) leptin receptors informing the brain of overall adiposity levels. In turn, the brain triggers compensatory changes to counter the increases in body fat, such as decreased energy intake and/or increased energy expenditure. Of course, if this hypothesized system functioned in this simplistic way, then no animals would become obese unless there are defects in the system. Indeed, defects do exist, as is most markedly seen in *ob/ob* and *db/db* mice that either do not synthesize leptin or do not have the long isoform leptin receptor, respectively (reviewed in Refs. 130–133). Regarding seasonal adiposity and Siberian hamsters, this hypothesized leptin feedback system, as typically conceived, is at odds with physiological reality. That is, although SD-exposed Siberian hamsters have decreased body fat and accordingly decreased WAT leptin mRNA (135–137) as well as circulating leptin concentrations (135, 138, 139), they have a concomitant decrease in food intake (13, 140, 141) rather than the predicted increase in food intake that would be expected from this leptin feedback model. Finally, the long isoform of the WAT leptin receptor gene expression also is reduced in the arcuate nucleus, a key integrative hypothalamic area for energy balance, in SDs compared with LDs (136), despite a predicted upregulation of the receptor according to many current notions regarding leptin and food intake.

The effects of exogenous leptin on body and lipid mass and food intake have also been examined in Siberian hamsters. Specifically, chronic leptin administration reverses the SD-induced decrease in food intake of Siberian hamsters to levels comparable with LD-housed controls not injected with leptin, but do not affect the food intake of their fatter LD-housed counterparts (142). This contrasts to the inhibition of food intake by leptin administration to genetically obese mice (*ob/ob* that are leptin-deficient; 143–146), but is consistent with decreased sensitivity to the suppression of

food intake by leptin of diet-induced obese AKR/J (147). In contrast to the above results in Siberian hamsters given leptin, another study report that leptin reduces body and lipid mass to a greater extent in SDs compared with LDs, but reduces food intake similarly in both photoperiods (135). The likely differences between these studies and those mentioned earlier may be due to differences in leptin administration that is critical for its effects on food intake and body fat (143, 148), as well as other methodological considerations (discussed in Ref. 142). Nevertheless, it appears that the effects of leptin on energy balance may be contingent upon the current photoperiod exposure. Therefore, seasonal changes in circulating leptin concentrations, coupled with changes in leptin sensitivity, may serve as part of an adaptive mechanism for increasing the odds of winter survival when food availability is decreased and adipose tissue stores are at their nadir.

**Summary of Possible Humeral Factors Mediating Levels of Seasonal Adiposity.** Finally, in terms of hormonal mediators of photoperiod/MEL-induced changes in obesity, it is possible that changes in several hormones working in combination are necessary to produce seasonal adiposity changes mediated by the photoperiod/MEL, and this may explain the difficulty in ascribing the SD-induced changes in adiposity to a single hormone. The ability to mimic accurately the natural secretion profiles for multiples of these hormones is difficult or impossible. Alternatively, it may be that MEL signals affect WAT and BAT via its innervation through the sympathetic nervous system (SNS) outflow from brain to fat and thereby modulate seasonal adiposity via this mechanism.

### The SNS May Modulate Seasonal Adiposity

WAT and BAT are both innervated by the SNS; (reviewed in Refs. 149–152). Whereas the innervation of BAT by the SNS has been undisputed and known for over 30 years (153), the innervation of WAT by the SNS has been more controversial and, to date, no convincing evidence exists for parasympathetic innervation of WAT (reviewed in Ref. 152). Although the neuroanatomical evidence for the SNS innervation of WAT was suggested over 100 years ago (154), and functional evidence for this innervation has been accumulating for nearly 90 years (155), some controversy about this innervation has continued until recently. The postganglionic sympathetic innervation of WAT was shown directly using retrograde and anterograde tract-tracing methods recently (156). Briefly, relatively separate populations of postganglionic neurons within the sympathetic chain corresponding to the thoracic and lumbar sections of the spinal cord was revealed for the inguinal WAT (IWAT) and epididymal WAT (EWAT) pads of Siberian hamsters (156). This study was prompted by an earlier finding showing that lipid was mobilized non-uniformly from WAT pads of SD-exposed Siberian hamsters (157). Specifically, the

more internally located pads (i.e., EWAT and retroperitoneal WAT [RWAT]) show the greatest lipid mobilization after SD exposure, as reflected by their decreased mass, whereas the more externally located pads (i.e., IWAT) show the least lipid mobilization after SD exposure (76, 117, 157). The identification of relatively separate neuroanatomical pools of postganglionic SNS neurons innervating these pads supports a likely anatomical basis for this differential lipid depletion, perhaps due to differences in sympathetic drives. Indeed, norepinephrine turnover, an indicator of SNS drive (158), was greater in EWAT than in IWAT corresponding to the greater decrease in EWAT mass than IWAT mass after SD exposure in Siberian hamsters (156).

Although partially satisfying as a mechanism underlying SD-induced increases in lipid mobilization from WAT as well as the fat pad-specific differences in the degree of lipid mobilization by Siberian hamsters, the CNS origins of the SNS outflow from brain to WAT were unknown until recently. Fortunately, the ability to use viral tract tracers to define complete neural circuits within the same animal, developed through the pioneering studies of Card (159) and Loewy (160), made this possible. This was accomplished using an attenuated strain of the pseudorabies virus (PRV; Bartha's K strain) as a retrograde transneuronal viral tract tracer and applied by us to define the SNS outflow from brain to WAT and BAT in Siberian hamsters and laboratory rats (159–164). Briefly, the use of this virus as a trans-synaptic retrograde tract tracer is possible because the virus is taken up into neurons following binding to viral attachment protein molecules found on the surface of neuronal membranes at the site of initial injection, in this case, WAT and BAT. These protein surface molecules act as “viral receptors.” Neurons synapsing on the infected cells become exposed to relatively high concentrations of the virus particles that have been exocytosed. The virus particles are then taken up by synaptic contact. This process continues, causing an infection along a hierarchical chain of functionally connected neurons from the adipose tissues, in this case, to the farthest reaches of the forebrain (165, 166). The virus is visualized using standard immunocytochemical methods or more recently using viruses engineered to make green fluorescent protein (167–169).

Application of this viral technology to the study of the innervation of WAT and BAT has revealed more similarities than differences in the CNS origins of the SNS outflow from brain to these two adipose tissue types (161, 162). Specifically, infected cells occur at all levels of the neuroaxis for both adipose tissue types and include the sympathetic chain, the spinal cord (intermediolateral cell column and central autonomic nucleus), the brainstem (the classically defined sympathetic sites such as the lateral and gigantocellular reticular nuclei, caudal raphe area, C1 adrenaline and A5 noradrenaline cell regions, and nucleus of the solitary tract), the midbrain (central gray), and the forebrain (paraventricular nucleus [PVN], SCN, lateral and dorsomedial hypothalamic nuclei, medial preoptic area, arcuate

nucleus, bed nucleus of the stria terminalis, and lateral septum); for details of all involved structures and differences between WAT and BAT, see References 161 and 162. The lack of infections in the ventromedial hypothalamus (VMH) after PRV injections to either WAT or BAT is notable because of the dozens of physiological studies indirectly suggesting VMH-SNS-WAT and VMH-SNS-BAT circuits (reviewed in ref. 150). The PRV technique has also failed to provide neuroanatomical support for the notion of circuits involving the VMH in the SNS innervation of a host of tissues including the pancreas (170) and adrenal medulla (160, 171). Most likely, the reason for the discrepancy between the results of the non-neuroanatomical studies and those of the PRV neuroanatomical studies is that manipulations of the VMH were not confined to this nucleus, causing ancillary stimulation or damage to the fibers of passage from the PVN, retrochiasmatic area, and dorsomedial, arcuate, and periventricular nuclei to brainstem and spinal cord sites important for the sympathetic control of adipose and other tissues underlying the metabolic and motivational aspects of ingestive behavior (reviewed in Ref. 151). Thus, we conclude that the VMH is not part of the SNS outflow to the periphery generally (152), or WAT specifically (161, 162) and does not participate in the mobilization of WAT via the sympathetic innervation of this tissue. This is not to suggest, however, that the VMH is not involved in seasonal responses per se; rather, the direct effects of the VMH on body fat and energy metabolism via the SNS are not supported. The VMH does appear to play an important role in reproductive behavior in general and a role in the photoperiodic control of reproductive status specifically. In terms of the latter, lesions of the VMH in Syrian hamsters showing SD-induced gonadal regression trigger accelerated gonadal recrudescence compared with pinealectomized hamsters or sham-operated controls (172). This suggests that an intact VMH is essential for the continued maintenance of SD-induced reproductive inhibition (172). Perhaps the continued stimulation of the neurons within the VMH that possess MEL receptors in Syrian hamsters (47) is necessary for maintenance of SD physiology in these hamsters.

### **The SNS Is a Possible Mechanism Underlying Photoperiod-Induced Changes in Seasonal Adiposity**

Here, we posit a possible mechanism whereby seasonal changes in adiposity may be mediated via alterations in SNS outflow to WAT, BAT, and the adrenal medulla in Siberian hamsters specifically, and perhaps in Syrian hamsters and other animals showing photoperiod/MEL-induced changes in body fat, more generally. We believe that the SCN is especially suited to coordinate seasonal changes because it functions as a biological clock that generates circadian behavioral and physiological rhythms (173), it appears vital for the reception of MEL-encoded photoperiod signals (33, 34, 52, 53), and it is part of the SNS outflow to WAT, BAT

(161, 162) and the adrenal medulla (174), as well as sympathetic and parasympathetic connections to a variety of other peripheral tissues (174; reviewed in Ref. 55) as revealed using the PRV tract tracing technique.


In terms of seasonal obesity and involvement of the SNS and SCN, we recently sought to determine if the SCN neurons that are part of the sympathetic outflow to WAT and BAT also express the MEL<sub>1a</sub> receptor, the functional MEL receptor subtype for seasonal responses. This was accomplished by labeling the sympathetic outflow from brain to WAT (164) and BAT (C-K. Song and T. Bartness, unpublished data) in Siberian hamsters using the PRV as a retrograde trans-synaptic tract tracer injected into IWAT or interscapular BAT combined with labeling of brain MEL<sub>1a</sub> receptors using *in situ* hybridization. Here, we will discuss the findings for WAT only. The SCN contains the highest number of MEL<sub>1a</sub> + PRV neurons compared with the other brain areas exhibiting MEL<sub>1a</sub> receptor gene expression. There are also, however, substantial numbers of MEL<sub>1a</sub> + PRV colocalized cells in the PVN, zona incerta, and reuniens/xiphoid area of the thalamus, dorsomedial nucleus, and to a lesser degree in the periventricular area fiber system, anterior hypothalamus, perifornical area, and paraventricular nucleus of the thalamus (PVT). Although the role of many of these areas in lipid mobilization (the response seen in SD-housed Siberian hamsters) is unclear or unknown, the SCN has been shown to be involved in increases in lipid mobilization generally (175–177; reviewed in ref. 150), and increases in photoperiod-induced lipid mobilization more specifically (33, 34, 52). For example, coronal microknife cuts just behind the SCN block the increases in lipid mobilization triggered by 24-hr fasts, forced exercise (swimming), cold exposure, and insulin-induced hypoglycemia in laboratory rats (178). Furthermore, microinfusions of SD-like MEL signals into the SCN trigger SD-like responses in Siberian hamsters (54), including decreases in body fat (25). Moreover, SCN lesions block the ability of SD-like exogenous MEL signals to decrease body and fat pad masses in pinealectomized Siberian hamsters (33, 34, 52, 150).

These data, strongly implicating the SCN in the SD-induced changes in body fat, added to what is known about the control of seasonal adiposity and other responses triggered by the changes in the photoperiod/MEL secretion duration, have been summarized graphically in Figure 1 (the number in brackets corresponds to the circled number in Fig. 1). Specifically, for photoperiod-induced body fat decreases in Siberian hamsters, we know the following.

The duration of night [1a] is the critical environmental stimulus triggering a decrease in body fat (i.e., with increasing duration in fall/winter; reviewed in Refs. 5 and 6).

This photic information is received by the retinal ganglion cells of the eye [1b] and transmitted to the SCN [2] via the retinohypothalamic tract and then to the pineal [3] through an identified multisynaptic pathway (16).

At the pineal [3], this photic information is transduced



**Figure 1.** Schematic illustration of the possible mechanism underlying the photoperiod-induced decreases in seasonal adiposity in Siberian hamsters. See text for a description of the system components and explanation for the numbers.

into a biochemical signal through the lengthening of the duration of MEL secretion that is roughly proportional to the lengthening of the dark period (10, 25).

Circulating MEL binds to the functional receptors mediating photoperiodic responses ( $MEL_{1a}$  receptors), especially those found on SCN neurons [4], but also elsewhere (51, 179), some of which are part of the CNS origins of the SNS outflow [5] to the periphery (164).

The stimulation of these  $MEL_{1a}$  receptors located on SNS outflow neurons by MEL increases the sympathetic drive on several peripheral tissues, including WAT [6a], BAT [6c], and adrenal medulla [6b].

Increases in the sympathetic drive (increased norepinephrine release; 180) via the SNS innervation of BAT [6a] (162, 181) increases BAT thermogenesis, growth, and/or thermogenic capacity [7] (180, 182).

Increases in the sympathetic drive on the adrenal medulla [6b] increases epinephrine release into the circulation,

affecting many tissues and promoting lipolysis in WAT [8] (183).

Increases in the sympathetic drive on WAT (156, 161) via its SNS innervation increases norepinephrine release from the nerve endings [6c] (156), also promoting WAT lipolysis [8] (183, 184).

Although the above hypothesized scenario focuses on the SNS outflow from brain to WAT as the primary mechanism underlying seasonal changes in adiposity by SD-housed Siberian hamsters, other factors are involved specifically, surgical (184) or chemical (183) sympathetic denervation of WAT only partially blocks SD-induced body fat decreases in these hamsters. Similarly, removal of only the adrenal medulla, the other branch of the sympathetic outflow, eliminates epinephrine-induced lipolysis, but only partially blocks SD-induced decreases in body fat (183). Sympathetic denervation of WAT combined with adrenal demedullation, however, completely blocks SD-induced

body fat decreases (183). Perhaps the ability of SCN lesions to completely block SD-induced lipid mobilization by Siberian hamsters is due to a centrally induced blockade of sympathetic outflow to both the adrenal medulla and WAT. It is unknown whether there are MEL receptors on neurons comprising the brain-SNS-adrenal medulla circuitry; however, the patterns of infection after PRV injections into the adrenal medulla of rats (171) or Siberian hamsters (M. Bamschad and T. Bartness, unpublished data) are quite similar to those seen after injection of PRV into WAT (161, 163). In addition, it may be that increases in energy expenditure via increased BAT thermogenesis and potential thermogenesis from pockets of ectopically induced brown adipocytes in WAT (185), as well as increased sensitivity to catecholamines in WAT, contribute to the decreases in seasonal adiposity in Siberian hamsters. That is, SD-exposed Siberian hamsters have increases in BAT norepinephrine turnover (180), as well as increases in gene expression for uncoupling protein-1 (UCP-1), one of a family of uncoupling proteins that is primarily responsible for the thermogenic activity of BAT (186), and peroxisome proliferator-activated receptor gamma (PPAR $\gamma$ ) coactivator-1 (PGC-1) (134), a critical co-activator of UCP-1 (187). There also may be increased sensitivity to catecholamines by Siberian hamster WAT, based on SD-induced increases in WAT  $\beta_3$ -adrenergic receptor gene expression in these animals (188). Finally, SD-exposed Siberian hamsters also show recruitment of dormant brown adipocytes and/or the induction of transdifferentiation of white adipocytes into brown adipocytes in their retroperitoneal WAT depots, as evidenced by the SD-induced increase in UCP-1 gene expression in this tissue (188). This effect seems identical to that occurring within otherwise similar white adipocyte populations when the sympathetic drive on WAT increases, such as occurs with cold exposure or pharmacological stimulation of  $\beta_3$ -adrenergic receptors, the major postganglionic sympathetic adrenergic receptor of WAT that, when stimulated, promotes lipolysis (185). Taken together, the SD-induced decreased seasonal adiposity of Siberian hamsters, and perhaps other species showing decreases in body fat in SDs, may be promoted by a coordinated suite of changes in WAT and BAT gene transcription that ultimately facilitates lipid mobilization and utilization of lipid fuels largely, but not exclusively (i.e., adrenal medullary catecholamines), due to increases in sympathetic drive to adipose tissues (see discussion in the "Summary" below).

The above speculation seems reasonable as a possible, or even likely, mechanism by which Siberian hamsters and other animals that get thinner in SDs decrease their lipid stores; however, what mechanisms underlie the SD-induced increase in adiposity in species such as Syrian hamsters? A parsimonious view of this phenomenon might be to assume that in Syrian hamsters and other species showing SD-induced body fat increases, MEL inhibits rather than stimulates the SNS drive on WAT. Support for this notion is indirect relative to WAT, but suggestive of an involvement

of the SNS. Specifically, SD exposure of Syrian hamsters decreases norepinephrine turnover in heart (189) with a concomitant LD-like norepinephrine turnover in BAT (107, 189, 190), despite increases in BAT mitochondrial content estimated by cytochrome c oxidase activity, and BAT mitochondrial proton conductance estimated by guanosine-5'-diphosphate (GDP) binding to isolated BAT mitochondria (190). In addition, it is possible that SDs reduce the sympathetic drive on WAT as in heart, although it has never been measured to date. This could not only promote lipid accumulation via a decrease in basal lipolysis levels, but also could increase adipocyte proliferation within the WAT pads. That is, decreases in sympathetic drive on WAT seen in many obesity models (191–193) including human obesity (1) is associated with increases in fat cell number (hyperplasia). This effect is most striking with elimination of sympathetic drive via denervation in hamsters (184) as well as rats (194). Indeed, SD-exposed Syrian hamsters have increases in fat cell number in several WAT pads (195). The SD-induced increases in adiposity are not primarily due to increases in food intake because the developing obesity often, but not always, occurs independently of significant overeating (11). Thus, it is possible that SDs decrease the sympathetic drive on WAT in these animals to promote increases in lipid storage while the sympathetic drive on BAT is unchanged, thereby promoting energy savings.

## Summary

We have presented data and speculation concerning the control of seasonal obesity by the photoperiod/MEL on its targets, and have primarily focused on the findings from the most commonly studied species showing seasonal changes in adiposity—Siberian and Syrian hamsters. The data to date are not compelling for a direct effect of MEL on WAT and BAT, despite some recent data to the contrary. Thus far, none of the possible hormonal intermediaries for the effects of MEL on seasonal adiposity appear likely as a mechanism by which MEL affects the photoperiodic control of body fat levels indirectly. One possible hormonal candidate that seems to deserve more study is glucagon. We also provided evidence pointing toward the SNS as a likely mediator of the effects of MEL on SD-induced body fat decreases in Siberian hamsters through increases in sympathetic drive on WAT and BAT, and we speculated that decreases in the SNS drive to adipose tissues may underlie the photoperiod-induced seasonal increases in adiposity of species such as Syrian hamsters. Clearly, we need to deepen our understanding of seasonal adiposity, although, to our knowledge, this is the only environmentally induced change in body fat where the key elements of its external trigger have been identified and can be traced to and through their transduction into a physiological stimulus that ultimately affects identified responses of white adipocyte physiology and cellularity (164 and Fig. 1 with associated text above). Finally, the comparative physiological approach to the study of sea-

sonal adiposity seems highly likely to yield significant insights into the mechanisms underlying this phenomenon and was the guiding stimulus for the our explorations into the SNS innervation of WAT.

The authors thank Drs. Ruth Harris and Eva Lacy, and Robert Bowers for their comments on this review and especially Drs. George Wade and Bruce Goldman for their roles in shaping this area of investigation and the career of the senior author in the areas of obesity, photoperiodism and their interaction.

1. Bjorntorp P. Possible mechanisms relating fat distribution and metabolism. In: Bouchard C, Johnson FE, Eds. *Fat Distribution During Growth and Later Health Outcomes*. New York: Alan R. Liss, pp175–191, 1988.
2. Coleman DL. Diabetes and obesity: thrifty mutants? *Nutr Rev* 36:129–132, 1978.
3. Mrosovsky N, Faust IM. Cycles of body fat in hibernators. *Int J Obesity* 9:93–98, 1985.
4. Bronson FH, Heideman PD. Seasonal regulation of reproduction in mammals. In: Knobil E, Neill JD, Eds. *The Physiology of Reproduction*. New York: Raven Press, pp541–583, 1994.
5. Bartness TJ, Wade GN. Photoperiodic control of seasonal body weight cycles in hamsters. *Neurosci Biobehav Rev* 9:599–612, 1985.
6. Bartness TJ, Fine JB. Melatonin and seasonal changes in body fat. In: Watson RR, Ed. *Melatonin in the Promotion of Health*. Boca Raton, FL: CRC Press, pp115–135, 1999.
7. Bjorntorp P. Adipose tissue distribution and function. *Int J Obesity* 15:67–81, 1991.
8. Himms-Hagen J. Brown adipose tissue metabolism and thermogenesis. *Annu Rev Nutr* 5:69–94, 1985.
9. Turek FW, Campbell CS. Photoperiodic regulation of neuroendocrine-gonadal activity. *Biol Reprod* 20:32–50, 1979.
10. Bartness TJ, Goldman BD. Mammalian pineal melatonin: a clock for all seasons. *Experientia* 45:939–945, 1989.
11. Bartness TJ, Wade GN. Photoperiodic control of body weight and energy metabolism in Syrian hamsters (*Mesocricetus auratus*): role of pineal gland melatonin gonads and diet. *Endocrinology* 114:492–498, 1984.
12. Hoffmann K. The influence of photoperiod and melatonin on testis size body weight and pelage colour in the Djungarian hamster (*Phodopus sungorus*). *J Comp Physiol* 85:267–282, 1973.
13. Wade GN, Bartness TJ. Effects of photoperiod and gonadectomy on food intake body weight and body composition in Siberian hamsters. *Am J Physiol* 246:R26–R30, 1984.
14. Kriegsfeld LJ, Nelson RJ. Gonadal and photoperiodic influences on body mass regulation in adult male and female prairie voles. *Am J Physiol* 270:R1013–R1018, 1996.
15. Dark J, Zucker I, Wade GN. Photoperiodic regulation of body mass food intake and reproduction in meadow voles. *Am J Physiol* 245:R334–R338, 1983.
16. Larsen PJ, Enquist LW, Card JP. Characterization of the multisynaptic neuronal control of the rat pineal gland using viral transneuronal tracing. *Eur J Neurosci* 10:128–145, 1998.
17. Moore RY. Neural control of the pineal gland. *Behav Brain Res* 73:125–130, 1996.
18. Vitale PM, Darrow JM, Duncan MJ, Shustak CA, Goldman BD. Effects of photoperiod pinealectomy and castration on body weight and daily torpor in Djungarian hamsters (*Phodopus sungorus*). *J Endocrinol* 106:367–375, 1985.
19. Hoffman RA. Seasonal growth and development and the influence of the eyes and pineal gland on body weight of golden hamsters (*M. Auratus*). *Growth* 47:109–121, 1983.
20. Bartness TJ, Goldman BD. Effects of melatonin on long day responses in short day-housed adult Siberian hamsters. *Am J Physiol* 255:R823–R830, 1988.
21. Bartness TJ, Goldman BD. Peak duration of serum melatonin and short day responses in adult Siberian hamsters. *Am J Physiol* 255:R812–R822, 1988.
22. Rhodes DH. The influence of multiple photoperiods and pinealectomy on gonads pelage and body weight in male meadow voles *Microtus pennsylvanicus*. *Comp Biochem Physiol A* 93:445–449, 1989.
23. Smale L, Dark J, Zucker I. Pineal and photoperiodic influences on fat deposition pelage and testicular activity in male meadow voles. *J Biol Rhythms* 3:349–355, 1988.
24. Hastings MH. Neuroendocrine rhythms. *Pharmacol Therapeutics* 50:35–71, 1991.
25. Bartness TJ, Powers JB, Hastings MH, Bittman EL, Goldman BD. The timed infusion paradigm for melatonin delivery: What has it taught us about the melatonin signal its reception and the photoperiodic control of seasonal responses? *J Pineal Res* 15:161–190, 1993.
26. Elliott JA, Goldman BD. Seasonal reproduction: photoperiodism and biological clocks. In: Adler NT, Ed. *Neuroendocrinology of Reproduction*. New York: Plenum Press, pp377–423, 1981.
27. Tamarkin L, Westrom WK, Hamill AI, Goldman BD. Effect of melatonin on the reproductive systems of male and female hamsters: a diurnal rhythm in sensitivity to melatonin. *Endocrinology* 99:1534–1541, 1976.
28. Wade GN, Bartness TJ. Seasonal obesity in Syrian hamsters: effects of age diet photoperiod and melatonin. *Am J Physiol* 247:R328–R334, 1984.
29. Bartness TJ, Wade GN. Body weight food intake and energy regulation in exercising and melatonin-treated Siberian hamsters (*Phodopus sungorus*). *Physiol Behav* 35:805–808, 1985.
30. Tamarkin L, Lefebvre NG, Hollister CW, Goldman BD. Effect of melatonin administered during the night on reproductive function in the Syrian hamster. *Endocrinology* 101:631–634, 1977.
31. Goldman BD, Hall V, Hollister C, Roychoudhury P, Tamarkin L. Effects of melatonin on the reproductive system in intact and pinealectomized male hamsters maintained under various photoperiods. *Endocrinology* 104:82–88, 1979.
32. Lynch GR, Hall ES. Effects of daily and chronic melatonin administration in *Peromyscus leucopus*: comparison with Syrian and Siberian hamsters. In: Brown GM, Wainwright SD, Eds. *The Pineal Gland: Endocrine Aspects*. New York: Pergamon Press, pp127–131, 1985.
33. Song CK, Bartness TJ. Microknife-cuts dorsal and caudal to the suprachiasmatic nucleus do not block short day responses by Siberian hamsters to timed infusions of melatonin. *Brain Res Bull* 45:239–246, 1998.
34. Song CK, Bartness TJ. The effects of anterior hypothalamic lesions on short-day responses in Siberian hamsters given timed melatonin infusions. *J Biol Rhythms* 11:14–26, 1996.
35. Bittman EL. The sites and consequences of melatonin binding in mammals. *Am Zool* 33:200–211, 1993.
36. Weaver DR, Rivkees SA, Carlson LL, Reppert SM. Localization of melatonin receptors in mammalian brain. In: Klein DC, Moore RY, Reppert SM, Eds. *Suprachiasmatic Nucleus: The Mind's Clock*. New York: Oxford University Press, pp289–308, 1991.
37. Morgan PJ, Barrett P, Howell E, Helliwell R. Melatonin receptors: localization molecular pharmacology and physiological significance. *Neurochem Int* 24:101–146, 1994.
38. Duncan MJ, Takahashi JS, Dubocovich ML. Characterization of 2-[125I]iodomelatonin binding sites in hamster brain. *Eur J Pharmacol* 132:333–334, 1986.
39. Weaver DR, Rivkees SA, Reppert SM. Localization and characterization of melatonin receptors in rodent brain by in vitro autoradiography. *J Neurosci* 9:2581–2590, 1989.
40. Acuna-Castroviejo D, Pabolos MI, Mendendez-Pelaez A, Reiter RJ. Melatonin receptors in purified cell nuclei of liver. *Res Commun Chem Pathol Pharmacol* 82:263–256, 1993.
41. Pang CS, Brown GM, Tang PL, Cheng KM, Pang SF. 2-[<sup>125</sup>I]iodomelatonin binding sites in the lung and heart: a link between the photoperiod signal melatonin and the cardiopulmonary system. *Biol Signals* 2:228–236, 1993.
42. Capsoni S, Viswanathan M, De Oliveira AM, Saavedra JM. Characterization of melatonin receptors and signal transduction system in rat arteries forming the circle of Willis. *Endocrinology* 135:373–378, 1994.
43. Helliwell RJA, Howell HE, Lawson W, Barrett P, Morgan PJ. Autoradiographic anomaly in <sup>125</sup>I-melatonin binding revealed in ovine adrenal. *Mol Cell Endocrinol* 104:95–102, 1994.

44. LeGouic S, Atgie C, Viguerie-Bascands N, Hanoun N, Larrouy D, Ambid L, Raimbault S, Ricquier D, Delagrance P, Guardiola-Lemaitre B, Penicaud L, Casteilla L. Characterization of a melatonin binding site in Siberian hamster brown adipose tissue. *Eur J Pharmacol* **339**:271–278, 1997.
45. Bittman EL, Weaver DR. Distribution of melatonin receptors in neuroendocrine tissues of the ewe. *Biol Reprod* **43**:986–993, 1990.
46. Duncan MJ, Takahashi JS, Dubocovich ML. Characteristics and autoradiographic localization of 2-[125I]iodomelatonin binding sites in Djungarian hamster brain. *Endocrinology* **125**:1011–1018, 1989.
47. Williams LM, Morgan PJ, Hastings MH, Lawson W, Davidson G, Howell H. E. Melatonin receptor sites in the Syrian hamster brain and pituitary: localization and characterization using [125I]iodomelatonin. *J Neuroendocrinol* **1**:315–320, 1989.
48. Reppert SM, Weaver DR, Rivkees SA, Stopa EG. Putative melatonin receptors in a human biological clock. *Science* **242**:78–81, 1988.
49. Duncan MJ, Mead RA. Autoradiographic localization of binding sites for 2-[125I]iodomelatonin in the pars tuberalis of the western spotted skunk (*Spilogale putorius latifrons*). *Brain Res* **569**:152–155, 1992.
50. Weaver DR, Reppert SM. Melatonin receptors are present in the ferret pars tuberalis and pars distalis but not in brain. *Endocrinology* **127**:2607–2609, 1990.
51. Reppert SM, Weaver DR, Ebisawa T. Cloning and characterization of a mammalian melatonin receptor that mediates reproductive and circadian responses. *Neuron* **13**:1177–1185, 1994.
52. Bartness TJ, Goldman BD, Bittman EL. SCN lesions block responses to systemic melatonin infusions in Siberian hamsters. *Am J Physiol* **260**:R102–R112, 1991.
53. Maywood ES, Buttery RC, Vance GHS, Herbert J, Hastings MH. Gonadal responses of the male Syrian hamster to programmed infusions of melatonin are sensitive to signal duration and frequency but not to signal phase nor to lesions of the suprachiasmatic nuclei. *Biol Reprod* **43**:174–182, 1990.
54. Badura LL, Goldman BD. Central sites mediating reproductive responses to melatonin in juvenile male Siberian hamsters. *Brain Res* **598**:98–106, 1992.
55. Bartness TJ, Song CK, Demas GE. SCN efferents to peripheral tissues: implications for biological rhythms. *J Biol Rhythms* **16**:196–204, 2001.
56. Ng TB, Wong CM. Effects of pineal indoles and arginine vasotocin on lipolysis and lipogenesis in isolated adipocytes. *J Pineal Res* **3**:55–66, 1986.
57. Ziotopoulou M, Mantzoros CS, Hileman SM, Flier JS. Differential expression of hypothalamic neuropeptides in the early phase of diet-induced obesity in mice. *Am J Physiol* **279**:E838–E845, 2000.
58. Prunet-Marcassus B, Ambid L, Viguerie-Bascands N, Penicaud L, Casteilla L. Evidence for a direct effect of melatonin on mitochondrial genome expression of Siberian hamster brown adipocytes. *J Pineal Res* **30**:108–115, 2001.
59. Hoffman RA, Hester RJ, Towns C. Effect of light and temperature on the endocrine system of the golden hamster (*Mesocricetus auratus* Waterhouse). *Comp Biochem Physiol* **15**:525–535, 1965.
60. Canguilhem B, Vaultier J-P, Pevet P, Coumaros G, Masson-Pevet M, Bentz I. Photoperiodic regulation of body mass food intake hibernation and reproduction in intact and castrated European hamsters *Cricetus cricetus*. *J Comp Physiol A* **163**:549–557, 1988.
61. Figala J, Hoffmann K, Goldau G. Zur Jahresperiodik beim Dsungarischen Zwerghamster *Phodopus sungorus* Pallas. *Oecologia* **12**:89–118, 1973.
62. Nelson RJ, Frank D, Smale L, Willoughby SB. Photoperiod and temperature affect reproductive and nonreproductive functions in male prairie voles (*Microtus ochrogaster*). *Biol Reprod* **40**:481–485, 1989.
63. Hasler JF, Buhl AE, Banks EM. The influence of photoperiod on growth and sexual function in male and female collared lemmings. *J Reprod Fert* **46**:323–329, 1976.
64. Dark J, Johnston PG, Healy M, Zucker I. Latitude of origin influences photoperiodic control of reproduction of deer mice (*Peromyscus maniculatus*). *Biol Reprod* **28**:213–220, 1983.
65. Whitsett JM, Underwood H, Cherry J. Photoperiodic stimulation of pubertal development in male deer mice: involvement of the circadian system. *Biol Reprod* **28**:652–656, 1983.
66. Gower BA, Nagy TR, Stetson MH. Effect of photoperiod testosterone and estradiol on body mass bifid claw size and pelage color in collared lemmings. *Gen Comp Endocrinol* **93**:459–470, 1994.
67. Hoffman RA, Davidson K, Steinberg K. Influence of photoperiod and temperature on weight gain food consumption fat pads and thyroxine in male golden hamsters. *Growth* **46**:150–162, 1982.
68. Campbell CS, Tabor J, Davis JD. Small effect of brown adipose tissue and major effect of photoperiod on body weight in hamsters (*Mesocricetus auratus*). *Physiol Behav* **30**:349–352, 1983.
69. Flint WE. Die Zwerghamster der Palaarktischen Fauna, Wittenberg, FRG: A. Ziemsen-Verlag 1966.
70. Steinlechner S, Heldmaier G, Becker H. The seasonal cycle of body weight in the Djungarian hamster: photoperiodic control and the influence of starvation and melatonin. *Oecologia* **60**:401–405, 1983.
71. Dark J, Zucker I. Gonadal and photoperiodic control of seasonal body weight changes in male voles. *Am J Physiol* **247**:R84–R88, 1984.
72. Nelson RJ, Kita M, Blom JM, Rhyne-Grey J. Photoperiod influences the critical caloric intake necessary to maintain reproduction among male deer mice (*Peromyscus maniculatus*). *Biol Reprod* **46**:226–232, 1992.
73. Blank JL, Korytko AI, Freeman DA, Ruf TP. Role of gonadal steroid and inhibitory photoperiod in regulating body weight and food intake in deer mice (*Peromyscus maniculatus*). *Proc Soc Exp Biol Med* **206**:396–403, 1994.
74. Morin LP, Fleming A. Variations of food intake and body weight with estrous cycle ovariectomy and estradiol benzoate treatment in hamsters (*Mesocricetus auratus*). *J Comp Physiol Psychol* **92**:1–6, 1976.
75. Slusser WN, Wade GN. Testicular effects on food intake body weight and body composition in male hamsters. *Physiol Behav* **27**:637–640, 1981.
76. Bartness TJ. Photoperiod sex gonadal steroids and housing density affect body fat in hamsters. *Physiol Behav* **60**:517–529, 1996.
77. Blank JL, Freeman DA. Differential reproductive response to short photoperiod in deer mice: role of melatonin. *J Comp Physiol* **169**:501–506, 1991.
78. Dark J, Whaling CS, Zucker I. Androgens exert opposite effects on body mass of heavy and light meadow voles. *Horm Behav* **21**:471–477, 1987.
79. Canguilhem B, Masson-Pevet M, Pevet P, Bentz I. Endogenous photoperiodic and hormonal control of the body weight rhythm in the female European hamster *Cricetus cricetus*. *Comp Biochem Physiol* **101**:465–470, 1992.
80. Dark J, Zucker I. Photoperiodic regulation of body mass and fat reserves in the meadow vole. *Physiol Behav* **38**:851–854, 1986.
81. Bartness TJ, Elliott JA, Goldman BD. Control of torpor and body weight patterns by a seasonal timer in Siberian hamsters. *Am J Physiol* **257**:R142–R149, 1989.
82. Hoffmann K. Effect of castration on photoperiodically induced weight gain in the Djungarian hamster. *Naturwissenschaften* **65**:494, 1978.
83. Goldman BD, Matt KS, Roychoudhury P, Stetson MH. Prolactin release in golden hamsters: photoperiod and gonadal influences. *Biol Reprod* **24**:287–292, 1981.
84. Ortavant R, Bocquier F, Pelletier J, Ravault JP, Thimonier J, Voland-Nail P. Seasonality of reproduction in sheep and its control by photoperiod. *Aust J Biol Sci* **41**:69–85, 1988.
85. Agius L, Robinson AM, Girard JR, Williamson DH. Alterations in the rate of lipogenesis in vivo in maternal liver and adipose tissue on premature weaning of lactating rats. *Biochem J* **180**:689–692, 1979.
86. Smith RW. Effect of pregnancy lactation and involution on metabolism of glucose by rat parametrial adipose tissue. *J Dairy Res* **40**:353–360, 1973.
87. Borer KT, Kelch RP, Corley K. Hamster prolactin: physiological changes in blood and pituitary concentrations as measured by a homologous radioimmunoassay. *Neuroendocrinology* **35**:12–21, 1982.
88. Ben-Jonathan N, Arbogast LA, Hyde JF. Neuroendocrine regulation of prolactin release. *Prog Neurobiol* **33**:399–347, 1989.
89. Bartness TJ, Wade GN, Goldman BD. Are the short photoperiod decreases in serum prolactin responsible for the seasonal changes in energy balance in Syrian and Siberian hamsters? *J Exp Zool* **244**:437–454, 1987.
90. Whitten RD, Youngstrom TG, Bartness TJ. Hyperprolactinemia does not promote testicular recrudescence in photoregressed Siberian hamsters. *Physiol Behav* **54**:175–178, 1993.

91. Cincotta AH, Meier AH. Reduction of body fat stores by inhibition of prolactin secretion. *Experientia* **43**:416–417, 1985.
92. Cincotta AH, MacEachern TA, Meier AH. Bromocriptine redirects metabolism and prevents seasonal onset of obese hyperinsulinemic state in Syrian hamsters. *Am J Physiol* **264**:E285–E293, 1993.
93. Cincotta AH, Schiller BC, Meier AH. Bromocriptine inhibits the seasonally occurring obesity hyperinsulinemia insulin resistance and impaired glucose tolerance in the Syrian hamster *Mesocricetus auratus*. *Metabolism* **40**:639–644, 1991.
94. Cincotta AH, Meier AH. Bromocriptine inhibits in vivo free fatty acid oxidation and hepatic glucose output in seasonally obese hamsters (*Mesocricetus auratus*). *Metabolism* **44**:1349–1355, 1995.
95. Cincotta AH, Wilson JM, deSouza CJ, Meier AH. Properly timed injections of cortisol and prolactin produce long-term reductions in obesity hyperinsulinaemia and insulin resistance in the Syrian hamster (*Mesocricetus auratus*). *J Endocrinol* **120**:385–391, 1989.
96. Collins S, Kuhn CM, Petro AE, Swick AG, Chrnyk BA, Surwit RS. Role of leptin in fat regulation. *Nature* **380**:677, 1996.
97. Vriend J. Evidence for pineal gland modulation of the neuroendocrine-thyroid axis. *Neuroendocrinology* **36**:68–78, 1983.
98. Vriend JB, Richardson BA, Vaughan MK, Johnson LY, Reiter RJ. Effects of melatonin on thyroid physiology of female hamsters. *Neuroendocrinology* **35**:79–85, 1982.
99. Ottenweller JE, Tapp WN, Pittman DL, Natelson BH. Adrenal thyroid and testicular hormone rhythms in male golden hamsters on long and short days. *Am J Physiol* **253**:R321–R328, 1987.
100. Nagy TR, Gower BA, Stetson MH. Photoperiod effects on body mass body composition growth hormone and thyroid hormones in male collared lemmings (*Dicrostonyx groenlandicus*). *Can J Zool* **72**:1726–1734, 1994.
101. Nagy TR, Gower BA, Stetson MH. Endocrine correlates of seasonal body mass dynamics in the collared lemming (*Dicrostonyx groenlandicus*). *Am Zool* **35**:246–258, 1995.
102. Seidel A, Heldmaier G, Schulz F. Seasonal changes in circulating levels of thyroid hormones are not dependent on the age in Djungarian hamsters *Phodopus sungorus*. *Comp Biochem Physiol* **88A**:71–7387
103. Masuda A, Oishi T. Effects of photoperiod temperature and testosterone-treatment on plasma T<sub>3</sub> and T<sub>4</sub> levels in the Djungarian hamster *Phodopus sungorus*. *Experientia* **45**:102–103, 1989.
104. O'Jile JR, Bartness TJ. Effects of thyroxine on the photoperiodic control of energy balance and reproductive status in Siberian hamsters. *Physiol Behav* **52**:267–270, 1992.
105. Puig-Domingo M, Guerrero JM, Menéndez-Pelaez A, Reiter RJ. Melatonin specifically stimulates type-II thyroxine 5'-deiodination in brown adipose tissue of Syrian hamsters. *J Endocrinol* **122**:553–556, 1989.
106. Reiter RJ, Klaus S, Ebbinghaus C, Heldmaier G, Redlin U, Ricquier D, Vaughan MK, Steinlechner S. Inhibition of 5'-deiodination of thyroxine suppresses the cold-induced increase in brown adipose tissue messenger ribonucleic acid for mitochondrial uncoupling protein without influencing lipoprotein lipase activity. *Endocrinology* **126**:2550–2554, 1990.
107. Sigurdson SL, Himms-Hagen J. Control of norepinephrine turnover in brown adipose tissue of Syrian hamsters. *Am J Physiol* **254**:R960–R968, 1988.
108. Bray GA, York DA. Hypothalamic and genetic obesity in experimental animals: an autonomic and endocrine hypothesis. *J Physiol* **377**:1–13, 1979.
109. Ottenweller JE, Tapp WN, Burke JM, Natelson BH. Plasma cortisol and corticosterone concentrations in the golden hamster (*Mesocricetus auratus*). *Life Sci* **37**:1551–1558, 1985.
110. Williams KA. Role of glucocorticoids in the photoperiodic control of body weight body fat and food intake in Siberian hamsters. Master's Thesis, Georgia State University, 1994.
111. Nelson RJ, Fine JB, Demas GE, Moffatt CA. Photoperiod and population density interact to affect reproductive and immune function in male prairie voles. *Am J Physiol* **270**:R571–R577, 1996.
112. Fischman AJ, Kastin AJ, Graf MV, Moldow RL. Constant light and dark affect the circadian rhythm of the hypothalamic-pituitary-adrenal axis. *Neuroendocrinology* **47**:309–316, 1988.
113. Allen-Rowlands CF, Allen JP, Greer MA, Wilson M. Circadian rhythmicity of ACTH and corticosterone in the rat. *J Endocrinol Invest* **3**:371–377, 1980.
114. Veldhuis JD, Iranmanesh A, Lizarralde G, Johnson ML. Amplitude modulation of a burstlike mode of cortisol secretion subserves the circadian glucocorticoid rhythm. *Am J Physiol* **257**:E6–E14, 1989.
115. Bartness TJ, McGriff WR, Maharaj MP, Borer KT. Photoperiodic control of serum insulin: effects on energy balance and reproductive status in Siberian hamsters. *Fed Am Soc Exp Biol Abstr*, p46, 1991.
116. Feldman D, Loose D. Glucocorticoid receptors in adipose tissue. *Endocrinology* **100**:398–405, 1977.
117. Bartness TJ. Short day-induced depletion of lipid stores is fat pad- and gender-specific in Siberian hamsters. *Physiol Behav* **58**:539–550, 1995.
118. Gray GO, Leavitt WW. RU486 is not an antiprogesterin in the hamster. *J Steroid Biochem* **28**:493–497, 1987.
119. Langley SC, York DA. Effects of antiglucocorticoid RU 486 on development of obesity in obese fa/fa Zucker rats. *Am J Physiol* **259**:R539–R544, 1990.
120. Okada S, York DA, Bray GA. Mifepristone (RU 486) a blocker of type II glucocorticoid and progesterin receptors reverses a dietary form of obesity. *Am J Physiol* **262**:R1106–R1110, 1992.
121. Miceli MO, Post CA, Woshowska Z. Abdominal vagotomy blocks hyperphagia and body weight gain in Syrian hamsters exposed to short photoperiod. *Soc Neurosci Abstr* **522.4**, 1989.
122. Baskin DG, Latterman DF, Seeley RJ, Woods SC, Porte D Jr, Schwartz MW. Insulin and leptin: dual adiposity signals to the brain for the regulation of food intake and body weight. *Brain Res* **848**:114–123, 1999.
123. Bartness TJ, McGriff WR, Maharaj MP. Effect of diabetes and insulin on photoperiodic responses in Siberian hamsters. *Physiol Behav* **49**:613–620, 1991.
124. LeMagnen J. Metabolic and feeding patterns: role of sympathetic and parasympathetic efferent pathways. *J Auton Nerv Sys* **10**:325–335, 1984.
125. Billington CJ, Briggs JE, Link JG, Levine AS. Glucagon in physiological concentrations stimulates brown fat thermogenesis in vivo. *Am J Physiol* **261**:R501–R507, 1991.
126. Yahata T, Kuroshima A. Metabolic cold acclimation after repetitive intermittent cold exposure in rat. *Jpn J Physiol* **39**:215–228, 1989.
127. Iwanij V, Amos TM, Billington CJ. Identification and characterization of the glucagon receptor from adipose tissue. *Mol Cell Endocrinol* **101**:257–261, 1994.
128. Goodman HM. The pancreas and regulation of metabolism. In: Mountcastle VB, Ed. *Medical Physiology*. St. Louis: C.B. Mosby, pp1638–1673, 1980.
129. Ahima RS, Flier JS. Adipose tissue as an endocrine organ. *Trends Endocrinol Metab* **11**:327–332, 2000.
130. Harris RBS. Leptin: much more than a satiety signal. *Annu Rev Nutr* **20**:45–75, 2000.
131. Campfield LA, Smith FJ, Burn P. The OB protein (leptin) pathway: a link between adipose tissue mass and central neural networks. *Horm Metab Res* **28**:619–632, 1996.
132. Clarke IJ, Henry B, Iqbal J, Goding JW. Leptin and the regulation of food intake and the neuroendocrine axis in sheep. *Clin Exp Pharmacol Physiol* **28**:106–107, 2001.
133. Trayhurn P, Hoggard N, Mercer JG, Rayner DV. Leptin: fundamental aspects. *Int J Obesity* **23**:22–28, 1999.
134. Gavrilova O, Marcus-Samuels B, Reitman ML. Lack of responses to a  $\beta$ -adrenergic agonist in lipoatrophic A-ZIP/F-1 mice. *Diabetes* **49**:1910–1916, 2000.
135. Klingenspor M, Niggemann H, Heldmaier G. Modulation of leptin sensitivity by short photoperiod acclimation in the Djungarian hamster *Phodopus sungorus*. *J Comp Physiol B* **170**:37–43, 2000.
136. Mercer JG, Moar KM, Ross AW, Hoggard N, Morgan PJ. Photoperiod regulates arcuate nucleus POMC AGRP and leptin receptor mRNA in Siberian hamster hypothalamus. *Am J Physiol* **278**:R271–R281, 2000.
137. Klingenspor M, Dickopp A, Heldmaier G, Klaus S. Short photoperiod reduces leptin gene expression in white and brown adipose tissue of Djungarian hamsters. *FEBS Lett* **399**:290–294, 1997.
138. Drazen DL, Kriegsfeld LJ, Schneider JE, Nelson RJ. Leptin but not immune function is linked to reproductive responsiveness to photoperiod. *Am J Physiol* **278**:R1401–R1407, 2000.
139. Horton TH, Buxton-OM, Losee-Olson S, Turek FW. Twenty-four-hour profiles of serum leptin in Siberian and golden hamsters: photoperiodic and diurnal variations. *Horm Behav* **37**:388–398, 2000.

140. Bartness TJ, Morley JE, Levine AS. Photoperiod-peptide interactions in the energy intake of Siberian hamsters. *Peptides* 7:1079-1085, 1986.
141. Masuda A, Oishi T. Effects of photoperiod and temperature on body weight, food intake, food storage, and pelage color in Djungarian hamster. *J Exptl Zool* 248:133-139, 1988.
142. Drazen DL, Demas GE, Nelson RJ. Leptin effects on immune function and energy balance are photoperiod dependent in Siberian hamsters (*Phodopus sungorus*). *Endocrinology* 142:2768-2775, 2001.
143. Campfield LA, Smith FJ, Guisez Y, Devos R, Burn P. Recombinant mouse OB protein: evidence for a peripheral signal linking adiposity and central neural networks. *Science* 269:546-549, 1995.
144. Halaas JL, Gajiwala KS, Maffei M, Cohen SL, Chait BT, Rabinowitz D, Lallone RL, Burley SK, Friedman JM. Weight-reducing effects of the plasma protein encoded by the obese gene. *Science* 269:543-546, 1995.
145. Pellemounter MA, Cullen MJ, Baker MB, Hecht R, Winters D, Boone T, Collins F. Effects of the obese gene product on body weight regulation in ob/ob mice. *Science* 269:540-543, 1995.
146. Weigle DS, Bukowski T, Foster D, Holderman S, Kramer J, Lasser G, Lofton-Day C, Prunkard D, Raymond C, Kuijper J. Recombinant ob protein reduces feeding and body weight in the ob/ob mouse. *J Clin Invest* 96:2065-2070, 1995.
147. Halaas JL, Boozer C, Blair-West J, Fidathusein N, Denton DA, Friedman JM. Physiological response to long-term peripheral and central leptin infusion in lean and obese mice. *Proc Natl Acad Sci U S A* 94:8878-8883, 1997.
148. Harris RB, Mitchell TD, Yan X, Simpson JS, Redmann SM Jr. Metabolic responses to leptin in obese db/db mice are strain dependent. *Am J Physiol* 281:R115-R132, 2001.
149. Himms-Hagen J. Neural control of brown adipose tissue thermogenesis hypertrophy and atrophy. *Front Neuroendocrinol* 12:38-93, 1991.
150. Bartness TJ, Demas GE, Song CK. Central nervous system innervation of white adipose tissue. In: Klaus S, Ed. *Adipose Tissue*. Georgetown, TX: Landes Bioscience, pp201-229, 2001.
151. Bartness TJ, Song CK, Demas GE. Central nervous system innervation of brown adipose tissue. In: Klaus S, Ed. *Adipose Tissue*. Georgetown, TX: Landes Bioscience, pp162-200, 2001.
152. Bartness TJ, Bamshad M. Innervation of mammalian white adipose tissue: implications for the regulation of total body fat. *Am J Physiol* 275:R1399-R1411, 1998.
153. Wirsén C. Studies in lipid mobilization with special reference to morphological and histochemical aspects. *Acta Physiol Scand* 65:1-46, 1965.
154. Dogiel AS. Die sensiblen Nervenendigungen im Herzen und in den Blutgefassen der Säugethiere. *Arch mikr Anat* 52:44-70, 1898.
155. Mansfeld G, Muller F. Der Einfluss der Nervensystem auf die Mobilisierung von Fett. *Arch Physiol* 152:61-67, 1913.
156. Youngstrom TG, Bartness TJ. Catecholaminergic innervation of white adipose tissue in the Siberian hamster. *Am J Physiol* 268:R744-R751, 1995.
157. Bartness TJ, Hamilton JM, Wade GN, Goldman BD. Regional differences in fat pad responses to short days in Siberian hamsters. *Am J Physiol* 257:R1533-R1540, 1989.
158. Young JB, Landsberg L. Suppression of the sympathetic nervous system during fasting. *Science* 196:1473-1475, 1977.
159. Card JP, Rinaman L, Schwaber JS, Miselis RR, Whealy ME, Robbins ME, Enquist LW. Neurotropic properties of pseudorabies virus: uptake and transneuronal passage in the rat central nervous system. *J Neurosci* 10:1974-1994, 1990.
160. Strack AM, Sawyer WB, Hughes JH, Platt KB, Loewy AD. A general pattern of CNS innervation of the sympathetic outflow demonstrated by transneuronal pseudorabies viral infections. *Brain Res* 491:156-162, 1989.
161. Bamshad M, Aoki VT, Adkison MG, Warren WS, Bartness TJ. Central nervous system origins of the sympathetic nervous system outflow to white adipose tissue. *Am J Physiol* 275:R291-R299, 1998.
162. Bamshad M, Song CK, Bartness TJ. CNS origins of the sympathetic nervous system outflow to brown adipose tissue. *Am J Physiol* 276:R1569-R1578, 1999.
163. Shi H, Bartness TJ. Neurochemical phenotype of sympathetic nervous system outflow from brain to white fat. *Brain Res Bull* 54:375-385, 2001.
164. Song CK, Bartness TJ. CNS sympathetic outflow neurons to white fat that express melatonin receptors may mediate seasonal adiposity. *Am J Physiol* 281:R666-R672, 2001.
165. Enquist LW, Card JP. Pseudorabies virus: a tool for tracing neuronal connections. In: Lowenstein PR, Enquist LW, Eds. *Protocols for gene transfer in neuroscience: towards gene therapy of neurological disorders*. New York, NY: John Wiley & Sons, pp333-348, 1996.
166. Loewy AD. Viruses as transneuronal tracers for defining neural circuits. *Neurosci Biobehav Rev* 22:679-684, 1998.
167. Irnaten M, Neff RA, Wang J, Loewy AD, Mettenleiter TC, Mendelowitz D. Activity of cardiorespiratory networks revealed by trans-synaptic virus expressing GFP. *J Neurophysiol* 85:435-438, 2001.
168. Smith BN, Banfield BW, Smeraski CA, Wilcox CL, Dudek FE, Enquist LW, Pickard GE. Pseudorabies virus expressing enhanced green fluorescent protein: a tool for in vitro electrophysiological analysis of trans-synaptically labeled neurons in identified central nervous system circuits. *Proc Natl Acad Sci U S A* 97:9264-9269, 2000.
169. Kim JS, Enquist LW, Card JP. Circuit-specific coinfection of neurons in the rat central nervous system with two pseudorabies virus recombinants. *J Virol* 73:9521-9531, 1999.
170. Jansen ASP, Hoffman JL, Loewy AD. CNS sites involved in sympathetic and parasympathetic control of the pancreas: a viral tracing study. *Brain Res* 766:29-38, 1997.
171. Strack AM, Sawyer WB, Platt KB, Loewy AD. CNS cell groups regulating the sympathetic outflow to adrenal gland as revealed by transneuronal cell body labeling with pseudorabies virus. *Brain Res* 491:274-296, 1989.
172. Bae HH, Mangels RA, Cho BS, Dark J, Yellon SM, Zucker I. Ventromedial hypothalamic mediation of photoperiodic gonadal responses in male Syrian hamsters. *J Biol Rhythms* 14:391-401, 1999.
173. Rusak B, Zucker I. Neural regulation of circadian rhythms. *Physiol Rev* 59:449-526, 1979.
174. Ueyama T, Krout KE, Nguyen XV, Karpitskiy V, Kollert A, Mettenleiter TC, Loewy AD. Suprachiasmatic nucleus: a central autonomic clock. *Nat Neurosci* 12:1051-1053, 1999.
175. Yamamoto H, Nagai K, Nakagawa H. Bilateral lesions of the SCN abolish lipolytic and hyperphagic responses to 2DG. *Physiol Behav* 32:1017-1020, 1984.
176. Gross J, Migliorini RH. Further evidence for a central regulation of free fatty acid mobilization in the rat. *Am J Physiol* 232:E165-E171, 1977.
177. Teixeira VL, Antunes-Rodrigues J, Migliorini RH. Evidence for centers in the central nervous system that selectively regulate fat mobilization in the rat. *J Lipid Res* 14:672-677, 1973.
178. Coimbra CC, Migliorini RH. Evidence for a longitudinal pathway in rat hypothalamus that controls FFA mobilization. *Am J Physiol* 245:E332-E337, 1983.
179. Song CK, Bartness TJ, Petersen SL, Bittman EL. Co-expression of melatonin (MEL1a) receptor and arginine vasopressin mRNAs in the Siberian hamster suprachiasmatic nucleus. *J Neuroendocrinol* 12:627-634, 2000.
180. McElroy JF, Mason PW, Hamilton JM, Wade GN. Effects of diet and photoperiod on NE turnover and GDP binding in Siberian hamster brown adipose tissue. *Am J Physiol* 250:R383-R388, 1986.
181. Seydoux J, Tribollet ER, Bouillaud F. Effectiveness of surgical denervation of interscapular brown adipose tissue in the rat: further observations. In: Hales JRS, Ed. *Thermal Physiology*. New York: Raven Press, pp197-199, 1984.
182. Heldmaier G, Steinlechner S, Rafael J, Vsiansky P. Photoperiodic control and effects of melatonin on nonshivering thermogenesis and brown adipose tissue. *Science* 212:917-919, 1981.
183. Demas GE, Bartness TJ. Direct innervation of body fat and adrenal medullary catecholamines mediate photoperiodic changes in body fat. *Am J Physiol* 281:1499-1505, 2001.
184. Youngstrom TG, Bartness TJ. White adipose tissue sympathetic nervous system denervation increases fat pad mass and fat cell number. *Am J Physiol* 275:R1488-R1493, 1998.
185. Cousin B, Cinti S, Morrioni M, Raimbault S, Ricquier D, Penicaud L, Casteilla L. Occurrence of brown adipocytes in rat white adipose tissue: molecular and morphological characterization. *J Cell Sci* 103:931-942, 1992.
186. Lin CS, Klingenberg M. Isolation of the uncoupling protein from brown adipose tissue mitochondria. *FEBS Lett* 113:299-303, 1980.

187. Puigserver P, Wu Z, Park CW, Graves R, Wright M, Spiegelman BM. A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis. *Cell* **92**:829–839, 1998.
188. Demas GE, Bowers RR, Bartness TJ, Gettys TW. Photoperiodic regulation of gene expression in brown and white adipose tissue of Siberian hamsters (*Phodopus sungorus*). *Am J Physiol* **282**(1): R114–R121, 2002.
189. Viswanathan M, Hissa R, George JC. Suppression of sympathetic nervous system by short photoperiod and melatonin in the Syrian hamster. *Life Sci* **38**:73–79, 1986.
190. McElroy JF, Wade GN. Short photoperiod stimulates brown adipose tissue growth and thermogenesis but not norepinephrine turnover in Syrian hamsters. *Physiol Behav* **37**:307–311, 1986.
191. Cleary MP, Brasel JA, Greenwood MRC. Developmental changes in thymidine kinase DNA and fat cellularity in Zucker rats. *Am J Physiol* **236**:E508–E513, 1979.
192. DiGirolamo M, Fine JB, Tagra K, Rossmanith R. Qualitative regional differences in adipose tissue growth and cellularity in male Wistar rats fed ad libitum. *Am J Physiol* **274**:R1460–R1467, 1998.
193. Faust IM, Johnson PR, Stern JS, Hirsch J. Diet-induced adipocyte number increase in adult rats: a new model of obesity. *Am J Physiol* **235**:E279–E286, 1978.
194. Cousin B, Casteilla L, Lafontan M, Ambid L, Langin D, Berthault M-F, Penicaud L. Local sympathetic denervation of white adipose tissue in rats induces preadipocyte proliferation without noticeable changes in metabolism. *Endocrinology* **133**:2255–2262, 1993.
195. Plunkett SS, Fine JB, Bartness TJ. Photoperiod and gender affect adipose tissue growth and cellularity in juvenile Syrian hamsters. *Physiol Behav* **70**:1–9, 2000.