

It is therefore not necessary to assume the local formation of antibodies in order to account for the active sensitization of pulmonary tissues.

102 (1280)

The rôle of hepatic tissue in anaphylaxis.

By W. H. MANWARING and HAROLD E. CROWE.

[*From the Department of Bacteriology and Experimental Pathology, Leland Stanford Jr. University.*]

If 0.25 per cent. goat serum in 50 per cent. defibrinated normal blood is repeatedly perfused through the liver of a normal guinea pig, a slight reduction in the toxicity of the perfusion fluid is usually observed, on subsequent tests with isolated anaphylactic lungs. In no case, however, is the reduction in toxicity sufficient to prevent the anaphylactic reaction in these lungs.

If the liver of an anaphylactic guinea pig is similarly perfused, the perfusion fluid usually becomes almost completely non-toxic for anaphylactic lungs.

This reduction in toxicity is not accompanied by a measurable decrease in the amount of goat protein in the perfusion fluid, as determined by a subsequent titration with a specific precipitin.

103 (1281)

The food value of soy bean products.

By THOMAS B. OSBORNE and LAFAYETTE B. MENDEL.

[*From the Laboratory of the Connecticut Agricultural Experiment Station and the Sheffield Laboratory of Physiological Chemistry in Yale University, New Haven, Connecticut.*]

Soy beans fed as the sole source of protein, or as a supplement to corn gluten, are suitable for the nutrition of rats. They contain sufficient water-soluble vitamine to promote normal growth; for diets containing soy bean flour, butter fat, starch, and an artificial salt mixture have promoted growth as well as comparable rations containing natural protein-free milk. The pres-