

MINIREVIEW

Iodine Metabolism and Thyroid-Related Functions in Organisms Lacking Thyroid Follicles: Are Thyroid Hormones also Vitamins? (44098)

J. G. EALES¹

Department of Zoology, University of Manitoba, Winnipeg, Manitoba, R3T 2N2, Canada

Abstract. Thyroid-related functions in organisms devoid of follicular thyroid tissue have been reviewed. In the lamprey, a primitive vertebrate, the larva concentrates iodide and synthesizes thyroid hormones (TH) by iodoperoxidase (IP)-mediated iodination of a thyroglobulin (TG)-like molecule in a subpharyngeal afollicular endostyle. The endostyle is the thyroid homolog, and it reorganizes into a follicular thyroid at metamorphosis to the adult. Ascidians and amphioxus, invertebrate protochordate relatives of vertebrates, also concentrate iodide and synthesize TH in a subpharyngeal afollicular endostyle, but the endostyle never transforms to follicles. Ascidian plasma contains L-thyroxine and its more biologically active derivative 3,5,3'-triiodo-L-thyronine, and TH receptors exist, but TH effects are poorly understood. No other invertebrates possess an endostyle. Several invertebrates concentrate iodide at other sites and form protein-incorporated iodohistidines and iodotyrosines; however, *de novo* iodothyronine biosynthesis through IP-mediated TG iodination has not been established. Nevertheless, TH occur in invertebrates, and exogenous iodotyrosines or iodothyronines have effects on jellyfish, insects, and sea urchins. Furthermore, gut bacteria metabolize TH, and plants may synthesize TH by nonenzymatic oxidative iodination. Thus, TH occur in many organisms and, after ingestion and enteric absorption, can enter the food chain. Indeed, sea urchin larvae obtain TH required to induce metamorphosis from plant diatoms. Thyroid hormones can therefore have vitamin-like effects and, in conjunction with vitamin D₃ and possibly with other steroids, may be more aptly termed vitamones. Availability of exogenous TH has implications for models of invertebrate and vertebrate TH metabolism and iodine salvaging, and it may explain the prominent and probable ancestral role of peripheral mechanisms in regulating thyroidal status.

[P.S.E.B.M. 1997, Vol 214]

¹ To whom requests for reprints should be addressed at Department of Zoology, University of Manitoba, Winnipeg, Manitoba, R3T 2N2, Canada. Aspects of this study were supported by a grant-in-aid of research to the author from the Natural Sciences and Engineering Council of Canada.

[P.S.E.B.M. 1997, Vol. 214]

0037-9727/97/2144-0302\$10.50/0

Copyright © 1997 by the Society for Experimental Biology and Medicine




Figure 1. Structures of selected iodoamino acids. TIH, 1,2,5-triiodo-histidine; DIT, 3,5-diiodotyrosine; T₄, L-thyroxine. Various degradative pathways are shown for T₄. Two other commonly encountered iodoamino acids not shown above are MIT (monoiodotyrosine, which lacks the 5-iodine of DIT) and T₃ (3,5,3'-triiodo-L-thyronine, which is formed from T₄ by removing the 5' iodine).

Thyroid tissue has been identified in all vertebrate species examined to date. It is characterized by highly vascularized "spheroidal" follicles consisting of a layer of epithelial cells surrounding a lumen filled with proteinaceous colloid. The collective function of the follicles is to ensure secretion of thyroid hormones (TH), which regulate metabolism. The primary TH are the iodothyronines, L-thyroxine (T₄) (Fig. 1), and 3,5,3'-triiodo-L-thyronine (T₃) which lacks one outer-ring iodine. Effects of TH differ among vertebrates, but the follicular appearance is remarkably uniform, and TH of fish are identical in structure to those of mammals (1). Thyroid follicles are found only in vertebrates, but it has been recognized since the late 19th century that thyroidal function is not confined to vertebrates with follicular thyroid tissue (reviewed in Refs. 2–10). Furthermore, the TH and their actions have now been reported in invertebrate animals and plants, and bacteria metabolize TH. The purpose of this review is to describe thyroid-related functions in invertebrate animals (Table I) and other organisms lacking a follicular thyroid. This survey shows that the occurrence and actions of TH and related iodocompounds extend to many organisms, and that TH pass through the food chain. Thus the definition of a hormone as an endogenously produced specific chemical messenger may not strictly apply to the TH. This has implications for interpreting TH metabolic pathways and their regulation in both invertebrate and vertebrate animals.

Criteria for Thyroid Function

What evidence for thyroid-related function might one anticipate in organisms lacking follicular thyroid tissue? Based on the vertebrate model (reviewed in Ref. 11), there may be evidence of TH synthesis, TH metabolism, TH actions, and regulation of thyroidal status.

TH Synthesis. Iodide (I⁻) is required for TH synthesis but is scarce in many environments. Thus, if an organism is forming TH it may need to concentrate iodide, probably by a specific energy-dependent carrier-mediated pump comparable to that in the vertebrate thyroid (Fig. 2). There will also be a need to iodinate and couple tyrosines to form iodothyronines (TH). In the thyroid, these processes are achieved through synthesis of two key proteins, thyroglobulin (TG) and iodoperoxidase (IP). Thyroglobulin provides the appropriate stereochemistry for tyrosine (tyrosyl) iodination and iodotyrosyl coupling; IP activates iodine to accomplish iodination and couples the iodotyrosyls to form iodinated TG (TGI). One might therefore anticipate a TG/IP system, or some equivalent, in an organism synthesizing TH. However, nonenzymatic mechanisms of TH formation have been demonstrated *in vitro* (12), which could occur *in vivo* (13). Thus TG/IP may not be obligatory for TH synthesis.

Iodoperoxidase is associated with the thyroid cell membrane and catalyzes TG iodination extracellularly at the colloid-cell membrane interface. Since IP and TG are co-packaged by the Golgi into vesicles prior to their secretion into the colloid, intracellular TGI formation could also occur in these vesicles (11) (Fig. 2). Such a "short-circuit" biosynthesis (14), although observed in cultured isolated thyrocytes, is not considered important in mammals *in vivo* (15). However, it indicates the feasibility of TH formation without follicles. In the thyroid, the iodothyronines are formed in the TGI molecules as covalently incorporated amino acids. The iodothyronines are stored as TGI in the luminal colloid whose borders continually undergo endocytosis. Endocytotic vesicles fuse with lysosomes, and proteolytic digestion within resulting phagolysosome releases TH, which pass to the blood (Fig. 2). Therefore, an animal lacking follicles but synthesizing TH might possess extra- or intracellular proteases capable of hydrolyzing TGI.

TH Metabolism. If an organism is synthesizing TH, then TH should occur in body fluids. Because of the low TH solubility in aqueous media, most TH will likely be bound reversibly to circulating proteins. In vertebrates, TH undergo several enzymatic conversions, including deiodination (removal of iodines from outer or inner iodothyronine rings), conjugations with glucuronic acid and sulfate, and quantitatively minor pathways involving deamination, decarboxylation, and diphenyl-ether cleavage (Fig. 1). These various conversions are

Table I. Taxonomic List of the Invertebrate Animals Included in the Present Survey of Iodine Metabolism and Thyroid Function

Phylum	Subphylum	Class	Common name
Chordata ^a	Urochordata	Ascidia	Tunicates
		Larvacea	Sea squirts
		Thaliacea	
	Cephalochordata	Branchiostomata	Amphioxus
Hemichordata		Enteropneusta	Acorn worms
Echinodermata		Echinoidea	Sea urchins
Bryozoa		Gymnolaemata	
Arthropoda		Insecta	Insects
		Malacostraca	Shrimps
			Crayfish
			Amphipods
Annelida		Branchiopoda	Daphnia
		Polychaeta	Sea worms
		Oligochaeta	Earth worms
Mollusca		Gastropoda	Abalone, snails
		Pelecypoda	Clams, bivalves
Nemertea		Enopla	Nemertine worms
Cnidaria		Scyphozoa	Jellyfish
		Hydrozoa	Hydra
		Anthozoa	Corals
Porifera		Calcarea	Sponges
		Demospongia	
		Sclerospongia	
		Hexactinellida	

^a The Cephalochordata and the Urochordata are often referred to as protochordates, since they represent the most primitive form of chordate organization which probably evolved into the vertebrates.

involved in the extrathyroidally regulated activation and inactivation of TH, and are anticipated in organisms using TH as chemical regulators.

TH Actions. Some rapid TH effects may be due to TH interaction with membrane and mitochondrial receptors, independent of regulation of protein synthesis. However, in vertebrates, TH bind to specific nuclear receptors that are part of a large superfamily of proteins which also serve as receptors for steroids, vitamin D₃, and retinoic acid. The hormone-occupied receptors bind in pairs to adjacent short DNA sequences (thyroid response elements) and constitute transcription factors that regulate specific gene expression. If TH exert biological actions in an organism, these actions are likely mediated by TH-specific nuclear or membrane receptors.



Regulation of Thyroidal Status. Thyroidal status in vertebrates is regulated both centrally and peripherally. Central regulation involves hypothalamic control of pituitary secretion of thyroid-stimulating hormone (TSH), which in turn stimulates TH synthesis and secretion. There is also negative feedback of free or unbound plasma TH to inhibit TSH secretion. Thus, with central control, plasma TH is determined primarily by the set-point of the hypothalamic-pituitary axis to free TH. In contrast, peripheral regulation adjusts activities of various extrathyroidal enzymatic pathways involved in metabolic conversions of TH. Of particular significance

is outer-ring deiodination of T₄ to biologically active T₃, but conjugation and inner-ring deiodination pathways also play roles in inactivating and degrading TH. If TH exert effects in an organism, then central and/or peripheral regulation of the biologically active form of TH (probably but not necessarily T₃) is anticipated.

Survey of Thyroid-Related Functions in Organisms Lacking Thyroid Follicles

Agnatha (Larval Lampreys). The most primitive extant vertebrates are the agnathans, the jawless cyclostomes (lampreys and hagfishes). Parasitic adult lampreys have thyroid tissue resembling that of higher vertebrates, but follicles are absent in their nonparasitic larvae (ammocoetes). However, ammocoetes have an elongated subpharyngeal bilobed glandular sac (subpharyngeal gland or endostyle) originating from the pharyngeal floor between the first and fifth branchial clefts (Fig. 3A). The subpharyngeal gland connects with the pharynx through a narrow hypobranchial duct. Stemming from observations of Schneider in 1879 (16), the endostyle is now established as the thyroid homolog. At metamorphosis the hypobranchial duct closes, part of the endostyle undergoes histolysis or involutes, and the remainder differentiates into thyroid follicles (17–20). Several cell types occur in the endostyle (17, 21–24), and, despite difficulties in following their individual fates during metamorphosis, the consensus is that the

Anatomical homology with the thyroid does not necessarily mean that the endostyle synthesizes TH. Early studies showed no endostylar stable iodine accumulation (29, 30), but there was uptake of radioiodide (31), and autoradiography showed that mainly type 2 and 3 cells concentrated radioiodide (22, 32, 33). Most endostylar radioiodine was protein-incorporated, and its location correlated with presumed TG (22). The presence of TG has since been confirmed using light and electron microscopic immunocytochemistry in type 2c and 3 cells and some type 5 cells (34–36). Iodoperoxidase has also been identified histochemically in the endostyle (37). Certain other ammocoete tissues, particularly notochord (38–40), concentrate radioiodide. However, *in vitro* enzymatic hydrolysis indicates that only the endostyle contains labeled MIT, DIT, T₃, and T₄ (9, 32, 41–44). Radiolabeled T₄ and some T₃ occur in ammocoete plasma following radioiodide injection (9, 32). High concentrations of TH, particularly T₃, have



How are TH transferred from endostyle to blood? TG may be secreted into the endostyle lumen and hydrolyzed by a protease to liberate TH (39, 47). Free TH may then be reabsorbed by endostylar cells, or the secretions may pass through the hypobranchial duct into the intestine where further TGI hydrolysis may complete TH release prior to TH absorption from the gut lumen (25). In either case, the endostyle acts as an exocrine tissue. Electron microscope autoradiography indicates most endostylar TG iodination occurs extracellularly at the cell/colloid interface (26). However, intracellular iodination and hydrolysis of TGI may also occur, and the TH so formed may then be released into either the lumen or the blood.

Putative T₃ receptors have been reported in ammocoete liver (48), but TH actions in ammocoetes remain unresolved. Thyroid hormones induce amphibian (49, 50) and piscine (51) metamorphoses and might therefore be expected to induce ammocoete metamorphosis. However, as ammocoete metamorphosis commences, plasma T₄ and T₃ levels decrease (45, 46, 52), and do not increase to a climax as in amphibians and fish (49, 53). Furthermore, ammocoete metamorphosis cannot be induced by iodine, anterior pituitary preparations,

or TH treatments (29, 30, 54, 55), suggesting no role for elevated TH levels in promoting ammocoete metamorphosis.

Hoheisel and Sterba (56) found that potassium perchlorate (KClO_4) induced ammocoete metamorphosis. Suzuki (57) also showed that, at least in the larger ammocoetes, KClO_4 and other goitrogens induced a normal metamorphosis accompanied by a decrease in thyroidal function. The phenomenon has been documented in greater detail in Youson's laboratory (58, 59). One interpretation is that high plasma TH levels suppress metamorphosis. Thus, when plasma TH levels are lowered by KClO_4 , metamorphosis can proceed. However, independent (extrathyroidal) actions of KClO_4 on metamorphosis cannot be excluded (59).

The endostyle is not pituitary regulated. Neither its histological appearance (55, 60) nor its radioiodide uptake (44) are altered by hypophysectomy. The endostyle is also insensitive to mammalian TSH preparations (40, 55, 61). Some endostylar cells hypertrophy in the presence of goitrogens (62–65), but the goitrogen response is not pituitary mediated, as it occurs after hypophysectomy (60), and is not abolished by T₄ treatment (64, 65). Central pituitary control over the thyroid is also absent in adult lampreys (66). Thus, regulation of thyroidal status in both larval and adult lampreys may occur instead in peripheral tissues through extrathyroidal adjustments of TH metabolism (1, 66).

In summary, the ammocoete endostyle carries out the functions of follicular thyroid tissue most likely by exocrine release of TGI into the gut lumen. The TH are liberated by either intra- or extracellular protease action, and TH probably enter the circulation from the gut lumen. The TH roles are incompletely understood. Regulation of thyroidal status does not seem to involve a central hypothalamo-hypophysial-thyroid/endostyle axis and is probably by peripheral mechanisms.

Cephalochordata. The cephalochordata (amphioxus, Branchiostoma) are marine chordate invertebrates with a notochord homologous to that of vertebrates. Together with the urochordata (see below) they are often termed protochordates. Both larvae and adults have an endostyle (Fig. 3B). In 1914, Van Wijhe (67) termed the amphioxus endostyle the "glandale thyroidea." Despite early debate, the amphioxus and ammocoete endostyles are now considered homologous (3, 6). The amphioxus endostyle exhibits thyroid-like function. It contains certain cell types that accumulate radioiodide (68–71); it contains a TG material with iodination properties remarkably similar to those of mammalian TG (72, 73); IP activity has been established (37, 74) and TH synthesis demonstrated (9, 75, 76).

Of particular interest is the endostylar site of TH iodination (69). Electron microscopic autoradiography indicates that, shortly after radioiodide administration, the silver grains, representing newly synthesized labeled

material, occur in the lumen close to type 5 cells (70, 74). Later, the grains are associated with cellular structures of type 5 and 6 cells, which may have taken up iodinated materials from the lumen in a manner similar to vertebrate TGI endocytosis (70).

The metabolism, regulation, and action of TH have not yet been studied.

Urochordata. The ascidians (tunicates or sea-squirts) are marine invertebrate chordates with a notochord, but lacking any cranial formation. They are mainly sessile suspension feeders with an endostyle producing a mucous filter as part of their feeding mechanism (Fig. 3C). The endostyle concentrates radioiodide from seawater (77–88) and contains proteins related immunologically to TG (89, 90) and to IP (87, 91, 92). Endostylar iodine is utilized to form MIT, DIT, and T₄ (79, 81, 83, 84, 91). Iodination and TH biosynthesis are restricted mainly to type 7 cells (87). Indeed digestive and thyroidal functions are distinct, and tissues associated with these two functions are spatially separated (93). Most of the TH synthesized by the endostyle undergo exocrine secretion into the gut lumen. Secretions are then carried by mucus through the pharyngeal cavity and may be absorbed by caudal cells in the alimentary tract (74).

Radioiodine is also incorporated into MIT and DIT in the tunic, the outer covering of the tunicate. Some workers report that iodothyronines are made there (90, 94, 95), but others contend that tunic iodinations are "nonthyroidal" in nature, resembling scleroprotein iodination found at the surface of skeletal tissues in some other invertebrates (81, 82). Several unusual tunic iodinated *L*-tyrosine alkaloids have been identified using nuclear magnetic resonance in a colonial ascidian (96); being cytotoxic, the alkaloids may defend against predation.

Fredriksson *et al.* (97, 98) recorded immunoreactive T₄ in low levels in plasma (0.2 ng/ml) and in extracts of the pharynx, alimentary canal, and tunic (2.7–8.4 ng/g). Tissue and plasma immunoreactive T₃ correlated with the distribution of administered [¹²⁵I]T₃, but the T₃ concentrations were much lower than those for T₄. However, the relatively high T₃ levels and T₃/T₄ ratio in the alimentary tissue suggested T₄-to-T₃ conversion at that site; deiodination of T₄ to T₃ has been demonstrated *in vivo* (99). Putative nuclear receptors for T₃ were identified in the pharynx and alimentary canal ($K_d = 1 \times 10^{-10}$ M) (98), but the receptor abundance was low (maximal binding capacity: 2.0–4.2 fmole T₃/mg protein).

Information on biological actions of TH in ascidians is sparse and based on potentially pharmacological TH levels (10^{-5} – 10^{-7} M) (98). In the compound ascidian, *Periphora*, stolen elongation was enhanced by T₄ or T₃ treatment (100); in *Phallusia*, treatment with TH increased activity of polyphenol oxidase, an enzyme associated with growth and development (101); in

Ascidia, metamorphosis to the adult was accelerated by T_4 (102). This latter effect is of interest since TH also induce development and metamorphosis in vertebrates (49, 51). However, free-swimming larvae do not concentrate radioiodide (102), and if TH do influence ascidian development the source and regulation of TH production at the larval stage need to be determined.

In summary, although the biological actions of TH and their mechanism of control have not been established, the ascidians do produce and metabolize TH. Furthermore, despite considerable anatomical differences in the endostyles of lamprey ammocoetes, amphioxus, and ascidians, the site of iodine organification in each case occurs in cells on the lateral and dorsal endostylar regions. Thus, there is a common protothyroid region (70). This is also true in the most primitive ascidians, the Larvacea (appendicularians) (103, 104), which retain several larval features and are free swimming as adults. They are of evolutionary interest since, as swimming ascidians, they may represent a form ancestral to vertebrates.

Hemichordata. The enteropneusts, or acorn worms, were once allied to chordates based on presence of gill slits and a notochord. However, their notochord is not homologous to that of chordates, and they are no longer considered close chordate relatives. Early workers found autoradiographic evidence of radioiodide binding by surface epidermal and glandular tissues (105). Later radiochromatographic studies involving prolonged immersion in seawater containing radioiodide revealed MIT but no DIT or iodothyronines (106, 107). Thus, there is no evidence that hemichordates produce iodothyronines.

Echinodermata. Using three species of sea urchin, Chino *et al.* (108) established that TH promote, in a dose-dependent manner, the metamorphosis of the pluteus larva. Thyroxine was effective at 10^{-10} – 10^{-9} M; T_3 at 10^{-9} – 10^{-8} M; 3,3',5'-triiodo-*L*-thyronine (reverse $T_3 = rT_3$) at 10^{-8} – 10^{-7} M; T_3 propionate at 10^{-7} – 10^{-6} M. In contrast to TH effects on vertebrates, T_4 potency exceeded that of T_3 , and rT_3 was modestly effective.

High-performance liquid chromatography (HPLC) and radioimmunoassay (RIA) techniques showed that larval TH contents increased during metamorphosis. The T_4 content at the four-arm stage was 0.05 pg/ 10^4 larvae, increasing to 1.5 pg/ 10^4 larvae at the six-armed stage and 2.9 pg/ 10^4 larvae at the eight-armed stage. The changes in T_3 were much greater, increasing from 5 pg/ 10^4 larvae at the six-armed stage to 143 pg/ 10^4 larvae at the eight-armed stage. Surprisingly, the larval TH content depended on the planktonic unicellular algal food that was required for metamorphosis to proceed. Algal T_4 content was 0.8 pg/ 10^9 cells; T_3 content was 18 pg/ 10^9 cells.

Thiourea and PTU did not block metamorphosis, suggesting that larval TH synthesis is not required for metamorphosis. However, both $KClO_4$ and KSCN, competitors of iodide transport, retarded development at concentrations of 1–10 mM. This inhibition was not completely offset by T_4 (10 nM) treatment, indicating toxic effects of the inhibitors or a co-requirement for iodide in development.

Developmental progress of the crown-of-thorns starfish, *Acanthaster planci*, through its larval stages is also accelerated by T_4 in a dose-dependent manner (109).

In summary, TH are important in echinoderm development, and there is strong evidence in sea urchins that ingested unicellular algae are the primary TH source. Receptors related to TH receptors may occur in sea urchins (110).

Bryozoa. Radioiodide was concentrated from seawater in the whole body of *Bugula neritina* and *Shizoporella errata* by ouabain- and dinitrophenol-sensitive transport (111–113). Uptake was blocked by potassium thiocyanate (KSCN), a probable competitor of iodide transport, but was unaffected by perchlorate or thiourea. The presence of organic iodine derivatives has not been studied.

Arthropoda (Insects and Crustaceans). Following an early observation of radioiodide incorporation into the *Drosophila* cuticle during tanning (114), Limpel and Caseda (115) injected radioiodide into representative insects from several orders and reported labeled MIT in the hemolymph, labeled DIT and T_4 in the muscle, and unidentified organic radioiodine derivatives in other tissues. Monoiodohistidine was the main excreted iodocompound. They examined the tissue distribution of the labeled materials in the cockroach, *Periplaneta americana*. *L*-thyroxine, MIT, and DIT were identified in muscle, brain, nerve cord, and fat body; T_3 was not identified conclusively by their single-dimension paper chromatography. Tong and Chaikoff (116) essentially repeated the above study on *Periplaneta*, but used three different paper chromatographic systems. They identified MIT and DIT, and reported other labeled organic derivatives, but found no labeled T_3 or T_4 . They concluded that the T_4 reported earlier (115) was probably a radioiodinated scleroprotein derivative.

Total iodine was measured in insects from a prairie river in Manitoba, Canada, in the North American goiter belt (117). Levels ranged from 0.3 to 8.57 μ g iodine/g dry wt. Highest levels were found in Trichopteran cases, supporting the view that organically incorporated iodine (probably iodoamino acids) may be formed in insect scleroproteins even in areas where iodine is not particularly abundant.

Administered TH influence insect physiology. *In vitro* addition of T_4 (3 μ M) to fat bodies of *Hyalophora*

cercropia promoted release of diglycerides and fatty acids within 3 hr (118). *In vivo* injection of T_4 into larvae of the silk moth, *Bombyx mori*, increased protein synthesis, RNA synthesis, silk production, and modified heart function (119). Subsequent studies on the same species confirmed that T_4 alters protein the nucleic acid content of the testis (120), fat body (121), and ovary (122), and alters the glycogen content in the fat body of the female (123). The response was biphasic and maximal at 1 ng T_4 /g body wt. Single injections of different doses of T_4 (0.5–5 ng/g) decreased the glycogen content of the ovary during the larval stage and increased it during pupal and adult stages (123, 124). In the same species, T_4 acted in a dose-dependent manner (0.5–2 ng/g) to increase Na^+/K^+ and Mg^{++} ATPases in larval and pupal tissues (gonads and silk gland); doses of 0.25 ng/g were usually ineffective; in adults, only the Na^+/K^+ ATPase in testis was increased (125). Also in the same species, feeding T_4 to larvae increased their growth as well as their silk and egg production, possibly due to an increase in circulating ecdysteroids (126, 127). In another silk moth, *Antheraea mylitta*, T_4 (0.5–1 ng/g) increased protein and amino acid turnover in ovary, testis, and fat body, and modified hemolymph amino acid profiles (128–131).

In all the above studies, T_4 was the administered TH, and in all but one instance (118) T_4 was either fed or injected. However, TH also exert a rapid (5- to 7-min) *in vitro* action to decrease the volume of the eggs of the locust, *Locusta migratoria* (132). This TH action resembled that of juvenile hormone (JH). The actions of both TH and JH were blocked by the Na^+/K^+ ATPase inhibitor, ouabain, and by ethoxzolamide, which specifically blocks JH receptors. Furthermore, the rapid effect of both hormones precluded actions through modified protein synthesis and indicated involvement of a membrane receptor. TH may closely resemble the phenoxypheols, such as fenoxycarb, that act as JH agonists after binding to the JH receptor. T_4 was more effective than T_0 (thyronine with no substituted iodines) but two orders of magnitude less potent than T_3 , which exerted a detectable action at 1 nM or lower. The nature of the insect JH receptor has not been determined, but the above results suggest that it could belong to the protein superfamily that includes the TH receptors.

Ecdysone is another hormone important in insect development. The ecdysone receptor is also related to the TH receptors and is a member of this superfamily (133–135). Thus, TH effects on ecdysone function are also possible.

Few crustacea have been studied but there is evidence of iodine organification. The total iodine contents in the prairie freshwater amphipod, *Hyaella*, and the decapod crayfish, *Astacus*, were respectively 1.15 and 4.87 μ g iodine/g dry wt (117). Scleroproteins in the developing exoskeleton of *Daphnia* incorporate radioi-

odine (105). Analysis by RIA of a shrimp diet for fish indicated a T_4 content of 10–15 ng/g and a T_3 content of 28 ng/g (136). Both T_4 and T_3 , as well as T_4 -to- T_3 conversion, have been detected by RIA and HPLC in the brine shrimp, *Artemia* (Brown C, personal communication). There are no reports on the metabolism or actions of TH in crustaceans.

In summary, there is yet no proof that arthropods synthesize TG, IP, or iodothyronines, although they can organify iodine to form protein-incorporated iodotyrosines and iodohistidines. Nevertheless, their tissues may contain immunoreactive T_3 and T_4 . Insects can also respond to exogenous TH and possess receptors in the same protein superfamily as TH receptors.

Annelida. Marine polychaete worms bind ambient radioiodide mainly in the scleroproteins of the pharyngeal teeth, cuticle, parapodia, and chaetae (105, 137). In the polychaete, *Nereis diversicolor*, iodine is absorbed from seawater at about 5 ng/worm/day, resulting in a whole-body content of 23–30 μ g of iodine/g (138). There is extensive whole-body iodine loss, but 15%–30% binds at the base of chaetae where tanning occurs. The iodotyrosines, MIT and DIT, were identified by paper chromatography; other labeled compounds did not correspond to iodothyronines or their derivatives.

In the oligochaete, *Eisenia foetida*, immunohistochemical analysis of the nervous system revealed TG-like activity which was not co-located with cholinesterase activity (139). This precludes misidentification of immunoreactive TG as cholinesterase, which shows striking homology to a TG subunit (140, and see below). Light microscopic autoradiography of radioiodide-injected worms showed silver grains selectively concentrated in the central nerve cord, particularly in the cerebral ganglion (141). The main concentration was between the neurosecretory cells and the neuropile fibers in the zone of the presumptive plexiform neurohemal complex. Treatment with methylmercaptimidazole (MMI), an IP inhibitor, did not alter setal radioiodine content, but radioiodination in the nervous tissue was blocked supporting an IP-iodinating system there.

In summary, iodine organification and iodotyrosine synthesis occur in the few annelids studied. However, despite indications of neural TG-like materials and an IP requirement, there is no direct evidence of iodothyronine formation.

Mollusca. Ambient radioiodide is incorporated by scleroproteins of pelecypod molluscs (105, 137). T_3 and T_2 , but not T_4 , were identified by paper chromatography in the snail, *Planorbis* (142). Tong and Chaikoff (116) found that radioiodide administered to snails was incorporated into MIT and DIT and other unidentified compounds. These iodocompounds were analyzed in three paper chromatographic systems, but did not correspond to iodothyronines or their derivatives.

Even though gastropod molluscs may not synthesize TH, they may have the potential to metabolize TH. Commercially available β -glucuronidase (Sigma Chemical Co., St. Louis, MO) prepared from abalone (*Haliotis*) entrails or the snail, *Helix pomatia*, contained contaminant T_4 outer-ring deiodination activity that was inhibited by propylthiouracil and blocked by excess T_4 (DiStefano JJ III, personal communication). This implies functional significance for conversion of T_4 to T_3 in these molluscs.

Based on data available from very few species, at least some molluscs may synthesize iodotyrosines. Synthesis of TH *de novo* has not been established, but some molluscs may be able to metabolize TH. There have been no studies on TH receptors or TH effects in mollusca.

Nemertea. Two nemertean worms, *Lineus ruber* and *Amphiporus angulatus*, incorporated radioiodide from seawater into the surface mucous coating (143). Paper chromatography of mucous extracts revealed labeled MIT, DIT, and T_3 , and at least one other unidentified organic material; T_4 was not identified. *L. ruber* mucus had a higher tyrosine content and a higher content of organically bound radioiodine than *A. angulatus*. The validity of the paper chromatographic analysis has not been checked by other methods.

Cnidaria (Coelenterata). Corals accumulate considerable quantities of iodine, and research primarily on the Gorgonacea, Pennatulata, and Antipatharia taxa established the location and identity of the iodocompounds (reviewed in Refs. 144 and 145). Significant quantities of MIT and DIT are synthesized in the structural scleroproteins. Indeed, DIT was originally termed gorgonin, or gorgonoic acid, because it was first identified from the gorgonid corals (146). The tyrosine content correlates with the degree of iodination, and both are greater in younger, more actively growing corals. However, there has been no convincing demonstration of iodothyronine synthesis. Lack of iodotyrosine coupling probably reflects the physical separation of the fibers containing the iodinated scleroproteins. The scleroproteins of two hydroids lacking the extensive skeletal structure of the corals also incorporated iodide from seawater and formed iodotyrosines (147). No biological effects of iodocompounds on corals have been documented.

There has also been interest in iodine metabolism and effects of iodomaterials in Scyphozoan jellyfish. The free-swimming medusae bud by strobilation from scyphistomae of sessile polyps. The strobilation process requires iodide and is induced by iodide and certain iodocompounds (148–154). Iodide concentrates in the polyp (155). This is probably due to trapping by organification, since iodide uptake was not blocked by perchlorate (156). Iodide efficacy in inducing strobilation was reduced by MMI, which blocks iodide organification

(156). Thus, contrary to earlier reports, iodide itself is not the active material. Based on radioiodine incorporation, the most common identifiable iodocompounds have been MIT and DIT. Labeled T_4 was reported by Spangenberg (149, 150) and by Black and Webb (152), but not by Silverstone *et al.* (156).

The most consistent induction of strobilation is with MIT and DIT. T_4 effects are equivocal. For example, T_4 may induce strobilation (150), exert no effect (154), or at high doses inhibit statolith development (151). Radioiodomaterials other than MIT and DIT have been identified chromatographically and might also induce strobilation (156). Formation of iodocompounds was greatest under environmental conditions favoring strobilation (29°C, 10^{-7} M iodide) (156), suggesting a regulatory function for these materials.

What are the tissue levels of MIT and DIT, do MIT and DIT have receptors, and do MIT and DIT undergo metabolic conversions?

Porifera. The sponges, particularly sclerospongiae, contain a fibrous scleroprotein, spongin, which incorporates iodine. In some species, the iodotyrosine content attains 14% of the total amino acids, but iodothyronines have not been reported in significant amounts (reviewed in Refs. 144, 145, and 157).

Protista. T_3 binds specifically to the membrane of the ciliate protozoan, *Tetrahymena pyriformis*, but binding has not been linked to any TH action and is reported to lead to T_3 endocytosis (158).

Bacteria. Studies on thyroid-related functions in bacteria are confined to *Escherichia coli* and other gastrointestinal bacteria of mammals. There is no evidence that TH are made by bacteria or exert actions on bacteria. However, bacteria participate in TH enteric metabolism.

Bacteria in the large intestine enzymatically hydrolyze TH-glucuronide and TH-sulfate conjugates to liberate unconjugated TH and other metabolites (159–161). As conjugates, TH are biologically less active, are hydrophilic, and are poorly absorbed across the intestinal wall (162). However, unconjugated TH are absorbed and contribute to an enterohepatic cycling of the biliary-excreted TH and their derivatives (163). Thus, the deconjugative bacterial activity increases the enteric TH pool available for enterohepatic cycling (161). However, there is no evidence that deconjugative bacterial activity is regulated in accordance with the physiological needs of the host body.

Nevertheless, T_4 and particularly T_3 , but not MIT, DIT, or iodide, are transported into cultured *E. coli* and become bound to intracellular proteins (164). Radioiodide is generated in the presence of radioiodinated T_4 or T_3 , indicating iodide removal from the labeled outer iodothyronine ring (161 and references cited, 164). However, labeled T_3 is not a product, indicating subsequent deiodination of the inner unlabeled tyrosyl ring.

Bacteria also decarboxylate and deaminate TH (161 and references cited). The consequences to the bacteria in metabolizing either the conjugated or unconjugated TH are unclear.

Plants. Marine algae concentrate halogens, including iodide, from seawater. Iodide is important for growth, morphogenesis, and reproduction of certain marine algae (165). Marine algae also form iodotyrosines (166). Until recently, there was no convincing evidence to support either formation or presence of iodothyronines by plants. However, with HPLC and RIA, T_4 (0.78 pg/ 10^9 cells) and T_3 (18.4 pg/ 10^9 cells) have been measured in unicellular marine algae (108). The mechanism of algal T_3 and T_4 formation was not studied, but it probably differs from that involving TG/IP in the thyroid. Under the conditions of high radiant solar energy available to plants, direct iodotyrosine oxidation and coupling may occur.

Through the use of gas chromatography-mass spectrometry-selective ion monitoring and also time-of-flight mass spectrometry, it was found that seeds of the plant *Sinapis alba* and the axillary bulbils of the tiger lily, *Lilium tigrinum*, could produce T_4 , but not T_3 , by a presumed nonenzymatic process (166). In the presence of oxidizing agents, such as ascorbic acid, extracts from these plants could convert DIT to T_4 . It is of future interest to examine other plants to determine the prevalence of formation and occurrence of iodocompounds.

The TH roles in plants are poorly understood. Thyroxine exerts an auxin-like activity in cotton seedlings, competing with indole acetic acid (IAA) for receptor sites, which suggests a developmental role (167). Thyroxine was effective at a concentration two to three orders of magnitude below that for IAA.

However, the main actions of iodocompounds may not be on the plants themselves, but serve to protect plants against organisms that infect or consume them. For example, iodotyrosines in marine algae are toxic to bacteria and may defend against such pathogens (166). Plant iodotyrosines may be harmful to vertebrates; if present in significant levels in blood, MIT and DIT depress catecholamine synthesis by blocking tyrosine hydroxylase (168). Plant iodothyronines may also affect the metabolism of the consumer. For example, both T_4 and T_3 mimic JH action on insect ovaries (132) and may interfere with insect metabolism and development. Plant TH may also elevate the thyroidal status of vertebrate consumers. This possibility is unexplored, but conversely some plants produce substances that depress the thyroidal status of their consumers. For example, cabbages and rapeseed contain the progoitrin/myrosinase combination. Unless removed by heat treatment or selective plant breeding, myrosinase converts progoitrin to goitrin, a potent antithyroid agent for humans, livestock, and fish (169, 170).

In summary, there is evidence that plants form iodotyrosines and iodothyronines, but probably by chemical reactions different from the enzymatic steps present in chordate animals. The extent of TH occurrence in plants needs to be explored. Thyroid hormones may have effects on plants; they may also protect plants against either invertebrate or vertebrate herbivores.

Summary of Evidence for Thyroid-Related Functions in Achordate Invertebrates

Neither thyroid follicles nor endostyles are found in invertebrates lacking a notochord (achordata), but is there other evidence of thyroid-related functions? The main quests have been for sites of radioiodide accumulation and radiochromatographic evidence of iodotyrosine and iodothyronine biosynthesis based on *de novo* radioiodide incorporation. Iodotyrosine formation occurs in numerous achordata, from sponges to insects. Iodothyronine formation has also been reported in a few taxa including insects, molluscs, nemertean worms, and jellyfish. However, several of these claims are based on single-dimensional paper chromatography, unsubstantiated by more rigorous immunoreactive, HPLC, or spectroscopy analyses. At present, *de novo* synthesis of iodothyronines in any achordate has not been established with certainty.

TH synthesis in invertebrate protochordates (ascidians and amphioxus) and vertebrates depends on TG and IP. Thus, presence of TG-like molecules and IP activity would also provide evidence of thyroid-like function. Immunoreactive TG has been reported in an annelid brain, which also concentrates radioiodide, but iodothyronine formation was not studied. TG presence in annelid nervous tissue could reflect the presumed common ancestry of TG and acetylcholinesterase (ACE). The carboxyl terminal half of TG shows a striking homology in amino acid sequences with that of ACE (171–174), suggesting a common ancestry for the two peptides. However, the divergence of TG from ACE is reported to stem from genic duplication of ACE close to the time of chordate evolution (175), which would not account for TG presence in the phylogenetically more ancient annelids.

Few attempts have been made to measure TH in achordate invertebrates. Immunoreactive iodothyronines have been reported in tissue extracts of arthropods and echinoderms. Occurrence of TH receptors, pathways for TH metabolism, or TH regulation have not been studied. However, there are several convincing biological effects of administered iodotyrosines and iodothyronines on jellyfish, insects, and larval echinoderms. It is unlikely they all represent nonphysiological or pharmacological TH effects. Consequently, it remains to be explained why in animals that do not appear to synthesize TH one can observe the presence and/or biological actions of TH.

Endogenous and Exogenous (Vitamin) Sources of Thyroid Hormones

In models of thyroid function, the TH source is usually assumed to be exclusively endogenous, but exogenous TH sources also exist. For example, dietary diatoms seem to provide the TH required to promote metamorphosis of larval sea urchins. Thus, exogenous TH of plant or animal origin could account for the occurrence and biological activity of TH in those invertebrates lacking *de novo* TH synthesis. Furthermore, the existence of exogenous TH sources also has important implications for interpretation of vertebrate thyroid function.

Exogenous TH are certainly available to vertebrates. TH occur in milk of eutherian mammals (176, 177) and marsupials (178); thus, both exogenous and endogenous TH sources may be utilized during development. TH also cross the placenta and provide an exogenous TH supply for the mammalian fetus (179). In fish and birds, the eggs may provide a TH source independent of *de novo* synthesis (180, 181). However, the main source of exogenous TH in vertebrates is probably from food. Vertebrates can absorb TH from the gastrointestinal tract (161), permitting the routine oral administration of prescribed TH to human patients. Although many naturally ingested iodocompounds are covalently incorporated (TGI or other iodoproteins), they may be hydrolyzed by intestinal proteases to liberate free TH, which can then be absorbed. Indeed, ascidians, amphioxus, and ammocoetes may secrete TGI directly from the endostyle into the gastrointestinal tract. Thus, the gut is probably the ancestral route for endogenous TH delivery to the circulation (182).

Vertebrates may obtain TH from plant and invertebrate foods and certainly from ingestion of any vascularized vertebrate tissue. Some vertebrates will ingest higher loads of exogenous TH than others. These include parasites such as vampire bats, lampreys, or hagfish subsisting on vertebrate blood or body fluids; predators consuming whole vertebrates (e.g., piscivorous fish or birds); or scavengers that eat carcass remnants. Inevitable ingestion of the thyroid and the gall bladder by some of these animals will provide a particularly rich TH source. Few vertebrate diets have been analyzed for TH content. Recent analysis of fish diets prepared from vertebrate and invertebrate sources revealed appreciable immunoreactive TH levels (T_4 , 10–45.2 ng/g diet and T_3 28–114 ng/g diet) (136).

The potential contribution of dietary TH to the plasma pool has received little attention by mammalian researchers. This may reflect the predominance of studies on chow-fed laboratory rats, which in the wild might consume diets with a higher TH content. However, significant thyroid function persists even in laboratory rats after complete and prolonged thyroidectomy (183, 184),




Figure 4. A flow diagram depicting pathways for the metabolism of exogenous and endogenous iodomaterials through the gut, hepatic-portal system, liver, bile, and thyroid. The model depicts the potential contributions of dietary TH and other iodomaterials to either the plasma iodide or TH pools. It emphasizes the roles of the liver both as a peripheral thyrostat and as a salvager of iodine from plasma TH exceeding immediate physiological needs.

and a dietary source of TH has been suggested (185). The human thyroid system certainly responds to dietary TH. A striking example is the outbreak of thyrotoxicosis without true hyperthyroidism, which occurred in 1984 in Minnesota, South Dakota, and Iowa (186), the consequences of which may still be felt (187). This outbreak was traced to ground beef contaminated with thyroid tissue. Thyroid-contaminated pork sausage may also have caused thyrotoxicosis (188).

To what extent does the inevitable ingestion of TH *routinely* contribute to the plasma TH pool? Ingestion of significant TH will not necessarily lead to overt hyperthyroidism. Within limits, exogenous TH may contribute to the circulating TH pool and thereby spare use of endogenous TH. Furthermore, autoregulatory deiodinative mechanisms which accommodate unpredictable influxes of exogenous TH to the plasma pool exist in peripheral tissues of both mammals (reviewed in Ref. 11) and fish (1) (Fig. 4).

Invertebrates and vertebrates may therefore derive TH from both exogenous and endogenous sources. In this respect, TH resemble vitamin D₃, a cholesterol derivative obtainable by humans as a vitamin from either plant or animal dietary sources, but also capable of being

formed in the dermis from cholesterol by ultraviolet radiation. Indeed, several other similarities exist between TH and vitamin D₃. They both interact with closely related nuclear receptors, and regulation of blood levels of their active forms (T₃ and 1,25(OH)₂-vitamin D₃) and inactive forms (reverse T₃ and T₂ or 24,25(OH)₂D₃ and 1,24,25(OH)₃D₃) is achieved by balancing remarkably analogous systems of activating and inactivating enzymes in peripheral tissues (189). Furthermore, they both may be formed in plants through the photic energy captured from sunlight. Both vitamin D₃ and T₃ are important regulatory chemicals; just as vitamin D₃ does not fit perfectly the definition of a vitamin (since it may be made in the dermis), neither may T₃ fit the exact definition of a hormone (since it may be obtained from the diet).

Exogenous Thyroid Hormones and Sources of Iodide for Salvaging

Thyroid function depends on iodine sources, which may be limiting in certain environments. Iodine insufficiency (endemic goiter) occurs in humans, but rarely in freshwater or terrestrial animals in their natural habitats. Research on dietary iodine availability has focused primarily on total dietary iodine intake or on inorganic (iodide) supplementation. However, as discussed above, a variable and significant proportion of the dietary iodine may include iodine organically incorporated as plant, invertebrate, or vertebrate substances, such as iodinated proteins (scleroproteins and TGI) and conjugated and nonconjugated iodothyronines and iodotyrosines. There may be a selective advantage for the consumer to access iodine from these varied organic sources. Selection pressure may have favored evolution of enzymes that liberate iodine in a form and at an anatomical site maximizing eventual iodine use by the thyroid (Fig. 4).

Iodine salvaging from iodothyronines involves iodothyronine-specific deiodinases. The mammalian type I deiodinase, which is active in liver, strips iodines from both rings (Fig. 1) and favors iodothyronine substrates after at least partial inactivation by sulfate conjugation (190). Therefore the liver is strategically placed to salvage iodine from excess TH present in portal blood flowing from the gut, and prior to blood passage to systemic vessels (Fig. 4). To date, enteric absorption of TH has been assumed to involve primarily endogenously produced TH, but ingested exogenous TH will be equally prone to uptake from the gut lumen. Thus, enteric absorption of TH not only permits reuse of a large extracorporal store of recyclable TH (163) but also contributes to iodide salvaging of TH from both endogenous and exogenous sources. Whether absorbed TH are recycled as such or degraded to recapture iodide will depend on the exogenous TH load relative to the demands of the body for TH.

As emphasized above, ingesta may also contain iodotyrosines, iodohistidines, and other iodocompounds. It will be advantageous to deiodinate them as well, either by intracellular iodotyrosine/iodohistidine deiodinases or by enteric bacteria. Iodotyrosines are cleared rapidly from rat plasma (191), and potent iodotyrosine deiodinases occur in the liver and other extrathyroidal tissues of several mammals (192–194). It is of future interest to explore extrathyroidal enzyme systems for salvaging iodine from organic sources other than iodotyrosines or iodothyronines.

Evolution of the Thyroid System and the Regulation of Thyroidal Status

The earliest roles of TH may have been as plant protectant substances or as vitamins. With endostyle evolution in protochordate ancestors, there was presumably a selective advantage in supplementing exogenous dietary TH with endogenous TH synthesis. This presumably led to selection of a TG-like molecule derived from a redundant cholinesterase genic duplication and to selection of an IP system promoting TG iodination. When marine chordate ancestors colonized relatively iodine-poor freshwater habitats there may have been strong selection pressures favoring the organization of thyroid tissue into follicles, with their high TH and iodine storage capacities. However, despite the evolution of a thyroid system for endogenous TH production, use of exogenous TH may persist in extant vertebrates, and indeed this is commensurate with TH and iodine economy. Such an open model (Fig. 4) has implications for the evolution of the regulation of thyroidal status.

The main regulation of thyroidal status in higher vertebrates involves the central hypothalamo-hypophysial-thyroidal axis. In conjunction with negative feedback, this axis ensures appropriate thyroidal secretion of T₄ (and T₃). Central control is supplemented by peripheral mechanisms regulating the balance between the formation and degradation of biologically active T₃. Despite the focus of much recent research, the peripheral control mechanisms in mammals and higher vertebrates tend to be considered secondary in importance to central control mechanisms. However, in lampreys, the most primitive extant vertebrates, no central control through a hypothalamo-hypophysial axis has been found, indicating that vertebrate thyroid function preceded its control by the brain and pituitary. In chordate ancestors relying on exogenous TH sources, peripheral mechanisms may have represented, as they currently do for vitamin D₃, the main means of regulating TH availability to target cells. This may reflect the evolutionary appearance of peripheral mechanisms for controlling thyroidal status in lower vertebrates, such as lampreys, prior to appearance of central control mechanisms (1, 195). It may also explain the persistence in higher vertebrates of peripheral mechanisms that con-

tribute to the regulation of thyroïdal status—mechanisms lacking in most other vertebrate endocrine systems.

Conclusions

Among invertebrates, only those with notochords (protochordates) possess an endostyle homologous to the vertebrate thyroid. Other invertebrates concentrate iodide, but the iodocompounds synthesized are primarily iodotyrosines. Despite a lack of convincing evidence for iodothyronine synthesis, some invertebrates contain iodothyronines and respond to exogenous iodothyronines. However, at least some plants can synthesize iodothyronines and may be a source of TH to their consumers. Thus, TH may pass through the food chain, and TH should be viewed as vitamins as well as hormones. Indeed, TH, vitamin D₃, and possibly other steroids comprise a class of regulatory chemicals that are available from both exogenous and endogenous sources, and which might be more aptly termed vitamones. Availability of exogenous TH has implications for interpreting TH metabolism, iodine salvaging, and TH regulation. It may also explain the prominent and probably ancestral role of peripheral mechanisms in regulating thyroïdal status.

I wish to thank Drs. C. Brown, K. G. Davey, J. J. DiStefano, III, J. M. McCleese, D. S. MacKenzie, and T. J. Wiens, as well as K. R. Finnson, K. Kohel, T. Narayansingh, and an anonymous referee for their help.

1. Eales JG, Brown SB. Measurement and regulation of thyroïdal status in teleost fish. *Rev Fish Biol Fisher* **3**:299–347, 1993.
2. Gorbman A. Some aspects of the comparative biochemistry of iodine utilization and the evolution of thyroïdal function. *Physiol Rev* **35**:336–346, 1955.
3. Barrington EJW. Some endocrinological aspects of the protochordata. In: Gorbman A, Ed. *Comparative Endocrinology*. New York: Wiley, pp250–282, 1959.
4. Berg O, Gorbman A, Kobayashi H. The thyroid hormones in invertebrates and lower vertebrates. In: Gorbman A, Ed. *Comparative Endocrinology*. New York: Wiley, pp302–309, 1959.
5. Gorbman A. Problems in the comparative morphology and physiology of the vertebrate thyroid gland. In: Gorbman A, Ed. *Comparative Endocrinology*. New York: Wiley, pp266–282, 1959.
6. Barrington EJW. Hormones and vertebrate evolution. *Experientia* **18**:201–210, 1962.
7. Olsson R. Endostyles and endostylar secretions. *Acta Zool* **44**:299–328, 1963.
8. Olsson R. General review of the endocrinology of the protochordata and Myxinoidea. *Gen Comp Endocrinol (Suppl 2)*:485–499, 1969.
9. Salvatore G. Thyroid hormone biosynthesis in agnatha and protochordata. *Gen Comp Endocrinol (Suppl 2)*:535–552, 1969.
10. Barrington EJW. Phylogenetic perspectives in vertebrate endocrinology. In: Barrington EJW, Ed. *Hormones and Evolution*. New York: Academic press, Vol I:pp1–46, 1979.
11. McNabb FMA. *Thyroid Hormones*. Englewood Cliffs, NJ: Prentice-Hall, 1992.
12. Cahnmann HJ, Funakoshi K. Model reactions for the biosynthesis of thyroxine. Nonenzymic formation of 3,5,3'-triiodothyronine from 4-hydroxy-3-iodophenylpyruvic acid, 3,5-diiodotyrosine, and oxygen. *Biochemistry* **9**:90–98, 1969.
13. Ito S, Sano K, Nishikawa M. Production of diiodohydroxyphenylpyruvic acid by nonenzymatic reaction. *Endocrinol Jpn* **16**:129–132, 1969.
14. Kuliawat R, Arvan P. Intracellular iodination of thyroglobulin in filter-polarized thyrocytes leads to the synthesis and basolateral secretion of thyroid hormone. *J Biol Chem* **269**:4922–4927, 1994.
15. Ekholm R. Iodination of thyroglobulin—An intracellular or extracellular process? *Mol Cell Endocrinol* **24**:141–163, 1981.
16. Schneider A. *Beitrage zur Vergleichenden Anatomie, und Entwicklungsgeschichte der Wirbelthiere*. Berlin: pp1–111, 1879.
17. Marine D. The metamorphosis of the endostyle of *Ammocoetes branchialis*. *J. Exp Med* **17**:379–395, 1913.
18. Kraentzel E. Contribution a l'etude de la Lamproie fluviatile, *Lampetra (Petromyzon) fluviatilis* L. I *Arch Biol (Paris)* **44**:469–517, 1933.
19. Leach JW. The endostyle and thyroid gland of the brook lamprey *Ichthyomyzon fossor*. *J Morphol* **65**:549–605, 1939.
20. Leach JW. The archetypal position of amphioxus and ammocoetes and the role of the endostyles in chordate evolution. *Am Nat* **78**:341–357, 1944.
21. Sterba G. Die Physiologie und Histogenese der Schilddruse und des Thymus beim Bachneunaugen (*Lampetra planeri* Bl.). *Wiss Z Friedrich-Schiller-Univ Jena* **2**:239–298, 1953.
22. Barrington EJW, Franchi LL. Some cytological characteristics of thyroïdal function in the endostyle of the ammocoete larva. *Q J Microscop Sci* **97**:393–409, 1956.
23. Egeberg J. Iodine-concentrating cells in the endostyle of ammocoetes. *Z Zellforsch* **68**:102–115, 1965.
24. Fujita H, Honma Y. Some observations on the fine structure of the endostyle of larval lampreys, ammocoetes of *Lampetra japonica*. *Gen Comp Endocrinol* **11**:111–131, 1968.
25. Clements-Merlini M, Gorbman A. The secretory cycle of the iodoproteins in ammocoetes. II. A radioautographic study of the transforming larval thyroid gland. *J Morphol* **106**:357–364, 1960.
26. Fujita H, Honma Y. Iodine metabolism of the endostyle of larval lampreys, ammocoetes of *Lampetra japonica*. *Z Zellforsch* **98**:525–537, 1969.
27. Wright G, Youson JH. Transformation of the endostyle of the anadromous sea lamprey, *Petromyzon marinus* L., during metamorphosis. I. Light microscopy and autoradiography with ¹²⁵I. *Gen Comp Endocrinol* **30**:243–257, 1976.
28. Wright G, Filosa MF, Youson JH. Immunocytological localization of thyroglobulin in the transforming endostyle of anadromous sea lampreys, *Petromyzon marinus* L., during metamorphosis. *Gen Comp Endocrinol* **42**:187–194, 1980.
29. Remy P. L'iode et la metamorphose de l'*Ammocoetes branchialis* et *Petromyzon planeri* Bloch. *C R Soc Biol Paris* **86**:129–131, 1922.
30. Horton FM. On the relation of the thyroid gland to metamorphosis in the lamprey. *J Exp Biol* **11**:257–261, 1934.
31. Gorbman A, Creaser CW. Accumulation of radioactive iodine by the endostyle of larval lampreys and the problem of the homology of the thyroid. *J Exp Zool* **89**:391–401, 1942.
32. Leloup J. Metabolisme de l'iode et fonctionnement endostylaire chez l'ammocoete de *Lampetra planeri*. *J Physiol* **47**:671–677, 1955.
33. Olivereau M. Presence d'iode lie organiquement dans certaines cellules endostylaires et dans la thyroïde. *Bull Assoc Anat* **92**:1113–1132, 1955.
34. Wright G, Filosa MF, Youson JH. Immunocytological localization of thyroglobulin in the endostyle of the anadromous sea lamprey, *Petromyzon marinus* L. *Am J Anat* **152**:263–268, 1978.
35. Suzuki S, Kondo Y. Thyroïdal morphogenesis and biosynthesis of thyroglobulin before and after metamorphosis in the lamprey, *Lampetra reissneri*. *Gen Comp Endocrinol* **21**:451–460, 1973.

36. Monaco F, Andreoli M, La Fosta A, Roche J. Thyroglobulin biosynthesis in a larval (ammocoete) and adult freshwater lamprey (*Lampetra planeri* Bl.). *Comp Biochem Physiol B* **60**:87–91, 1978.
37. Tsuneki K, Kobayashi H, Ouji M. Histochemical distribution of peroxidase in amphioxus and cyclostomes with special reference to the endostyle. *Gen Comp Endocrinol* **50**:189–200, 1983.
38. Leloup J, Fontaine M. Iodine metabolism in lower vertebrates. *Ann N Y Acad Sci* **86**:316–353, 1960.
39. Clements-Merlini M, Gorbman A. The secretory cycle of the iodoproteins in ammocoetes. I. A radioautographic time study of the subpharyngeal gland. *J Morphol* **106**:337–356, 1960.
40. Clements-Merlini M. Metabolism of I^{131} by the endostyle and notochord of ammocoetes under different conditions of temperature. *Gen Comp Endocrinol* **2**:240–248, 1962.
41. Leloup J, Berg O. Sur le presence d'acides amines iodes (monoiodotyrosine, diiodotyrosine, et thyroxine) dans l'endostyle de l'ammocoete. *C R Soc Biol Paris* **238**:1069–1071, 1954.
42. Salvatore G, Covelli I, Sena L, Roche J. Fonction thyroïdienne et métabolisme de l'iode chez la larve d'un cyclostome (*Petromyzon planeri* Bl. *C R Soc Biol Paris* **153**:1686–1692, 1959.
43. Roche J, Salvatore G, Covelli I. Métabolisme de l'iode et fonction thyroïdienne chez la larve (Ammocoete) d'un Cyclostome. *Petromyzon planeri* Bl. *Comp Biochem Physiol* **2**:90–99, 1961.
44. Pickering AD. Effects of hypophysectomy on the activity of the endostyle and thyroid gland in the larval and adult river lamprey, *Lampetra fluviatilis* L. *Gen Comp Endocrinol* **18**:335–343, 1972.
45. Wright G, Youson JH. Serum thyroxine concentrations in larval and metamorphosing anadromous sea lamprey, *Petromyzon marinus* L. *J Exp Zool* **202**:27–32, 1977.
46. Lintlop SP, Youson JH. Concentration of triiodothyronine in sera of the sea lamprey, *Petromyzon marinus*, and the brook lamprey, *Lampetra lamottenii*, at various stages of their life cycle. *Gen Comp Endocrinol* **49**:187–194, 1983.
47. Clements M, Gorbman A. Protease in ammocoetes endostyle. *Biol Bull* **108**:258–263, 1955.
48. Lintlop SP, Youson JH. Binding of triiodothyronine to hepatocyte nuclei from sea lampreys *Petromyzon marinus* at various stages of the life cycle. *Gen Comp Endocrinol* **49**:428–436, 1983.
49. Galton VA. Thyroid hormone action in amphibian metamorphosis. In: Oppenheimer JH, Samuels HH, Eds. *Molecular Basis of Hormone Action*. New York: Academic Press, pp445–483, 1983.
50. Galton VA. The role of thyroid hormones in amphibian development. *Am Zool* **28**:309–318, 1988.
51. Inui Y, Miwa S. Thyroid hormone induces metamorphosis of flounder larvae. *Gen Comp Endocrinol* **60**:450–454, 1985.
52. Leatherland JF, Hilliard RW, Macey DJ, Potter IC. Changes in serum thyroxine and triiodothyronine concentrations during metamorphosis of the southern hemisphere lamprey *Geotria australis*, and the effect of propylthiouracil, triiodothyronine and environmental temperature on serum thyroid hormone concentrations of ammocoetes. *Fish Physiol Biochem* **8**:167–177, 1990.
53. Miwa S, Tagawa M, Inui Y, Hirano T. Thyroxine surge in metamorphosing flounder larvae. *Gen Comp Endocrinol* **70**:158–163, 1988.
54. Young JZ, Bellerby CW. The response of the lamprey to injection of anterior lobe pituitary extract. *J Exp Biol* **12**:246–252, 1935.
55. Knowles FGW. The duration of the larval life in ammocoetes and an attempt to accelerate metamorphosis by injections of an anterior-pituitary extract. *Proc Zool Soc London* **111**:101–109, 1941.
56. Hoheisel LG, Sterba G. Über die Wirkung von Kaliumperchlorat ($KClO_4$) auf Ammocoeten von *Lampetra planeri* Bloch. *Z Mikrosk Anat Forsch* **70**:490–516, 1963.
57. Suzuki S. Induction of metamorphosis and thyroid function in the larval lamprey. In: Mederios-Neto G, Gaitan E, Eds. *Frontiers in Thyroidology*. New York: Plenum Press, pp667–670, 1986.
58. Holmes J, Youson JH. Induction of metamorphosis in landlocked sea lampreys, *Petromyzon marinus*. *J Exp Zool* **267**:598–604, 1993.
59. Youson JH. Environmental and hormonal clues and endocrine glands during lamprey metamorphosis. In: Davey KG, Peter RE, Tobe SS, Eds. *Perspectives in Comparative Endocrinology*. National Research Council of Canada, Ottawa, pp400–407, 1994.
60. Barrington EJW, Sage M. On the response of the endostyle of the hypophysectomized larval lamprey to thiourea. *Gen Comp Endocrinol* **7**:463–474, 1966.
61. Olivereau M. Endostyle de l'ammocoete (*Lampetra planeri* Bloch) et hormone thyroïdienne. *C R Assoc Anat* **43**:636–657, 1956.
62. Jones RP. Effect of thiourea on the endostyle of ammocoetes. *Nature* **160**:638, 1947.
63. Klenner J, Schipper AL. The response of the endostyle of *Lampetra lamottenii* to iturmil, thiouracil, thiourea, thiocyanate and thyroxine. *Anat Rec* **120**:790, 1954.
64. Barrington EJW, Sage M. On the responses of the glandular tracts and associated regions of the endostyle of the larval lamprey to goitrogens and thyroxine. *Gen Comp Endocrinol* **3**:153–165, 1963.
65. Barrington EJW, Sage M. On the responses of the iodine-binding regions of the endostyle of the larval lamprey to goitrogens and thyroxine. *Gen Comp Endocrinol* **3**:669–679, 1963.
66. Dickhoff WW, Darling DS. Evolution of thyroid function and its control in lower vertebrates. *Am Zool* **23**:697–707, 1983.
67. Wijhe v JW. Studien über Amphioxus. I. Mund und Darmkanal während der Metamorphose. *Verh Kon Akad Wetensch (2 Sect)* **18**(1). Amsterdam: pp1–84, 1914.
68. Thomas IM. The accumulation of radioactive iodine by *Amphioxus*. *J Mar Biol Ass U K* **35**:203–210, 1956.
69. Barrington EJW. The localization of organically bound iodine in the endostyle of *Amphioxus*. *J Mar Biol Ass U K* **35**:117–119, 1958.
70. Fredriksson G, Ericson LE, Olsson R. Iodine binding in the endostyle of *Branchiostoma lanceolatum* (Cephalochordata). *Gen Comp Endocrinol* **56**:177–184, 1984.
71. Ericson LE, Fredriksson G, Ofverholm T. Ultrastructural localization of the iodination centre in the endostyle of the adult amphioxus (*Branchiostoma lanceolatum*). *Cell Tissue Res* **241**:267–273, 1985.
72. Dominici R, Monaco F, Andreoli M, De Pirro R, Demetrio G, Roche J. Presence de thyroglobuline et biosynthèse des hormones thyroïdiennes dans l'endostyle de l'Amphioxus (Cephalochorde), et evolution. *C R Soc Biol Paris* **175**:68–71, 1981.
73. Monaco F, Dominici R, Andreoli M, De Pirro R, Roche J. Thyroid hormone formation in thyroglobulin synthesized in the amphioxus (*Branchiostoma lanceolatum*). *Comp Biochem Physiol B* **70**:341–344, 1981.
74. Fredriksson G, Ofverholm T, Ericson LE. Electron-microscopic studies of iodine-binding and peroxidase activity in the endostyle of the larval amphioxus (*Branchiostoma lanceolatus*). *Cell Tissue Res* **241**:257–266, 1985.
75. Covelli I, Salvatore G, Sena L, Roche J. Sur la formation d'hormones thyroïdiennes et de leur précurseurs par *Branchiostoma lanceolatus* Pallas (Amphioxus). *C R Soc Biol Paris* **154**:1165–1169, 1960.
76. Tong W, Kerkoff P, Chaikoff IL. Identification of labelled thyroxine and triiodothyronine in Amphioxus treated with ^{131}I . *Biochem Biophys Acta* **56**:326–331, 1962.
77. Gorbman A. Identity of an iodine-storing tissue in ascidian. *Science* **94**:102, 1941.
78. Barrington EJW. The distribution and significance of organically bound iodine in the ascidian *Ciona intestinalis*. *J Mar Biol Ass U K* **36**:1–16, 1957.

79. Roche J, Salvatore G, Rametta G, Varrone S. Sur la presence d'hormone thyroïdiennes (3:5:3'-triiodothyronine et thyroxine) chez un Tunicien (*Ciona intestinalis* L.). C R Soc Biol Paris **153**:1751–1757, 1959.
80. Barrington EJW, Thorpe A. An autoradiographic study of the binding of ^{125}I in the endostyle and pharynx of the ascidian *Ciona intestinalis* L. Gen Comp Endocrinol **5**:373–385, 1965.
81. Barrington EJW, Thorpe A. The identification of monoiodotyrosine, diiodotyrosine and thyroxine in extracts of the endostyle of the ascidian, *Ciona intestinalis* L. Proc Roy Soc Ser B **163**:136–149, 1965.
82. Barrington EJW, Thorpe A. Histochemical and biochemical aspects of iodine binding in the tunic of the ascidian, *Dendrodoa grossularia* (van Beneden). Proc Roy Soc Ser B **171**:91–109, 1968.
83. Kennedy GR. The distribution and nature of iodine in ascidians. Gen Comp Endocrinol **7**:500–511, 1966.
84. Suzuki S, Kondo Y. Demonstration of thyroglobulin-like iodinated proteins in the branchial sac of tunicates. Gen Comp Endocrinol **17**:402–406, 1971.
85. Thorpe A, Thorndyke MC, Barrington EJW. Ultrastructural and histochemical features of the endostyle of the ascidian *Ciona intestinalis* with special reference to the distribution of bound iodine. Gen Comp Endocrinol **19**:559–571, 1972.
86. Thorpe A, Thorndyke MC. The endostyle in relation to iodine binding. Symp Zool Soc Lond **36**:159–177, 1979.
87. Dunn AD. Ultrastructural autoradiography and cytochemistry of iodine-binding cells in the ascidian endostyle. J Exp Zool **188**:103–124, 1974.
88. Fredriksson G, Ofverholm T, Ericson LE. Iodine binding and peroxidase activity in the endostyle of *Salpa fusiformis*, *Thalia democratica*, *Doliolotta gegenbauri*, and *Doliolum nationalis* (Tunicata, Thaliacea). Cell Tissue Res **253**:403–411, 1988.
89. Thorndyke M. Evidence for a mammalian thyroglobulin in the endostyle of the ascidian *Styela clava*. Nature **271**:51–52, 1978.
90. Dunn AD. Studies on iodoproteins and thyroid hormones in ascidians. Gen Comp Endocrinol **40**:473–483, 1980.
91. Dunn AD. Properties of an iodinating enzyme in the ascidian endostyle. Gen Comp Endocrinol **40**:484–493, 1980.
92. Fujita H, Sawano F. Fine structural localization of endogenous peroxidase in the endostyle of ascidians, *Ciona intestinalis*. A part of phylogenetic studies of the thyroid gland. Arch Histol Jpn **42**:319–326, 1979.
93. Fredriksson G, Ericson LE, Ofverholm T. The endostyle; A double organ. Am Zool **26**:108A, 1986.
94. Roche J, Salvatore G, Rametta G. Sur la presence et la biosynthese d'hormones thyroïdiennes chez un Tunicien, *Ciona intestinalis* L. Biochim Biophys Acta **63**:154–165, 1962.
95. Roche J, Rametta G, Varrone S. Metabolisme de l'iode et formation d'iodothyronine (T_3 et T_4) au cours de la regeneration de la tunique chez une ascidie, *Ciona intestinalis* L. Gen Comp Endocrinol **42**:277–284, 1964.
96. Carroll AR, Bowden BF, Coll JC. Studies of Australian ascidians. 2. Novel cytotoxic iodotyrosine-based alkaloids from colonial ascidians, *Alpidum* sp. Austral J Chem **46**:825–832, 1993.
97. Fredriksson G, Lebel JM, Leloup J. Thyroid hormones and putative T_3 receptors in a protochordate, *Phallusia mammillata* Cuvier. Gen Comp Endocrinol **82**:255–256, 1991.
98. Fredriksson G, Lebel JM, Leloup J. Thyroid hormones and putative nuclear T_3 receptors in tissues of the ascidian, *Phallusia mammillata* Cuvier. Gen Comp Endocrinol **92**:379–387, 1993.
99. Leloup J, Seugnet I. In vivo thyroxine monodeiodination in the ascidian, *Phallusia mamillata*. Gen Comp Endocrinol **74**:276–277, 1989.
100. Fukumoto M. Experimental control of budding and stolen elongation in *Perophora orientalis*, a compound ascidian. Dev Growth Res **13**:73–88, 1971.
101. Thorndyke MC. An *in vivo* Stimulating effect of 3,5',5-triiod-L-thyronine on polyphenol oxidase activity in an ascidian. J Endocrinol **58**:679–680, 1973.
102. Patricolo E, Ortolani G, Cascio A. The effect of thyroxine on the metamorphosis of *Ascidia malaca*. Cell Tissue Res **214**:289–301, 1981.
103. Olsson R. The cytology of the endostyle of *Oikopleura dioica*. Ann N Y Acad Sci **118**:1038–1051, 1965.
104. Fredriksson G, Ofverholm T, Ericson LE. Ultrastructural demonstration of iodine binding and peroxidase activity in the endostyle of *Oikopleura dioica* (Appendicularia). Gen Comp Endocrinol **58**:319–327, 1985.
105. Gorbman A, Clements M, O'Brien R. Utilization of radioactive iodine by invertebrates with special study of several annelida and mollusca. J Exp Zool **127**:75–92, 1954.
106. Roche J, Rametta G, Salvatore G. Sur la fixation de l'iode par un Enteropneuste (*Glossobalanus minutus* Kow.) et sa significance. C R Soc Biol Paris **150**:1902–1906, 1961.
107. Barrington EJW, Thorpe A. Comparative observations on iodine binding by *Saccoglossus horsti* Brambell and Goodhart, and by the tunic of *Ciona intestinalis* L. Gen Comp Endocrinol **3**:166–175, 1963.
108. Chino Y, Saito M, Yamasu K, Suyemitsu T, Ishihara K. Formation of the adult rudiment of sea urchins is influenced by thyroxine. Dev Biol **161**:1–11, 1994.
109. Johnson LG, Cartwright CM. Thyroxine-accelerated larval development in the crown-of-thorns starfish, *Acanthaster planci*. Biol Bull **190**:299–301, 1996.
110. Chan SM, Xu ND, Niemeyer CC, Bone JR, Flytzanis CN. SPCOUP-TF—A sea-urchin member of the steroid-thyroid hormone receptor family. Proc Natl Acad Sci U S A **89**:10568–10572, 1992.
111. Roche J, Andre S, Covelli I. Sur la fixation et l'utilisation de l'iode ^{131}I par deux bryozoaires, *Bugula neritina* L. et *Schizoporella errata* (Walters 1878). Comp Biochem Physiol **11**:215–221, 1964.
112. Andre S. Sur la biochimie comparee du transport des iodures. Decouplage des phosphorylations oxydatives par le 2,4-dinitrophenol et fixation de l'iode par un Bryozoaire (*Bugula neritina* L.) et une Algue (*Ulva* sp.), a la lumiere et a l'obscurite. C R Soc Biol Paris **158**:1465–1470, 1964.
113. Andre S. Sur la biochimie comparee du transport des iodures. Action de divers inhibiteurs sur la fixation des iodures par un Bryozoaire (*Bugula neritina* L.) et par deux Algues marines (*Laminaria flexicaulis* et *Enteromorpha* sp.). C R Soc Biol Paris **59**:2327–2332, 1965.
114. Wheeler BM. Halogen metabolism of *Drosophila gibberosa*—I. Iodine metabolism studied by means of ^{131}I . J Exp Zool **115**:83–104, 1950.
115. Limpel LE, Casada JE. Iodine metabolism in insects. I. *In vivo* metabolism of radionuclide. J Exp Zool **135**:19–27, 1957.
116. Tong W, Chaikoff IL. ^{131}I utilization by the aquarium snail and the cockroach. Biochim Biophys Acta **48**:347–351, 1961.
117. Gregory LA, Eales JG. Factors contributing to high levels of plasma iodide in brook trout, *Salvelinus fontinalis* (Mitchill). Can J Zool **53**:267–277, 1975.
118. Bhaktan NMG, Gilbert LI. Effect of some vertebrate hormones on lipid mobilization in the insect fat body. Gen Comp Endocrinol **11**:186–197, 1968.
119. Medda AK, Ray AK, Dasgupta AC, Rey CD. Thyroid hormone actions in magur fish and silk worms. In: Stockigt JR, Nagataki S, Eds. Thyroid Research VIII. Canberra, Australia: Australian Academy of Sciences, pp240–243, 1981.
120. Chaudhuri A, Medda AK. Changes in protein and nucleic acid contents of male gonad of silkworm, *Bombyx mori*, at different developmental stages after thyroxine treatment. Proc Nat Acad Sci India **56**:301–306, 1986.

121. Chaudhuri A, Medda AK. Thyroxine-induced alterations in protein and nucleic acid content of fat body of female silkworm, during different developmental stages. *Insect Sci Appl* **8**:43–48, 1987.
122. Chaudhuri A, Medda AK. Effect of thyroxine on protein, RNA and DNA content of ovary of silkworm, *Bombyx mori*, at larval, pupal, and adult stages of development and production of eggs. *Zool Jb Anat* **115**:85–90, 1987.
123. Chaudhuri A, Medda AK. Influence of thyroxine on ovarian glycogen of *Bombyx mori* L (race Nistari) during ontogeny. *Insect Sci Appl* **14**:621–626, 1994.
124. Chaudhuri A, Medda AK. Thyroxine-induced alterations in glycogen content of fat body of female silkworms, *Bombyx mori* (race Nistari), during larval, pupal and adult stages of development. *Ann Ent* **10**:17–21, 1992.
125. Reddy KD, Chaudhuri AB, Sukumar K. Enrichment of ion-specific adenosine-triphosphatase activities by thyroxine in different tissues of the silkworm, *Bombyx mori* L. during development. *Insect Biochem Mol Biol* **24**:243–248, 1994.
126. Thyagaraja BS, Kelly TJ, Masler EP, Borkovec AB. Thyroxine-induced hemolymph protein and ecdysteroid increases in the silkworm, *Bombyx mori*: Effect on larval growth and silk production. *J Insect Physiol* **37**:153–159, 1991.
127. Thyagaraja BS, Masler EP, Kelly TJ, Borkovec AB. Thyroxine-induced changes in ovarian protein and ecdysteroid levels in the silkworm, *Bombyx mori* L: Effect on ovarian maturation and egg production. *Comp Biochem physiol* **104A**:247–253, 1993.
128. Reddy KD, Chaudhuri A, Thangavelu K. L-thyroxine (T_4) elevates the free amino acid pool of hemolymph plasma of tasar silkworm, *Antheraea mylitta* Drury (Lepidoptera, Saturniidae). *Horm Metab Res* **26**:570–573, 1994.
129. Reddy KD, Chaudhuri A, Thangavelu K. Thyroxine effect on protein and nucleic acid turnover of testis and ovary during 5th larval stage of non-diapausing tasar silkworm, *Antheraea mylitta* (Saturniidae: Lepidoptera). *Indian J Exp Biol* **32**:409–412, 1994.
130. Reddy KD, Chaudhuri A, Thangavelu K. Effect of L-thyroxine-sodium pentahydrate on protein and nucleic acid turnover in fatbody during 5th larval stage of non-diapausing tasar silkworm, *Antheraea mylitta* (Saturniidae: Lepidoptera). *Indian J Exp Biol* **32**:413–417, 1994.
131. Reddy KD, Chaudhuri A, Thangavelu K. L-thyroxine (T_4) elevates the free amino acid pool of haemolymph plasma of tasar silkworm, *Antheraea mylitta* (Saturniidae: Lepidoptera). *Indian J Exp Biol* **32**:418–422, 1994.
132. Davey KG, Gordon DRB. Fenoxycarb and thyroid hormones have JH-like effects on the follicle cells of *Locusta migratoria* *in vitro*. *Arch Insect Biochem Physiol* **32**:613–622, 1996.
133. Hatzivassiliou E, Cardot P, Zannis V, Mitsialis A. A *Drosophila* RXR- α homolog can mobilize the thyroid-hormone receptor to activate the human apolipoprotein-A-II (APOA-II) promoter. *Circulation* **90**:402, 1994.
134. Horner MA, Chen TH, Thummel CS. Ecdysteroid regulation and DNA-binding properties of *Drosophila* nuclear hormone-receptor superfamily members. *Dev Biol* **168**:490–502, 1995.
135. Antoniewski C, Laval M, Lepesant JA. Structural features critical to the activity of an ecdysone receptor-binding site. *Insect Biochem Mol Biol* **23**:105–114, 1993.
136. MacKenzie DS, Moon HY, Gatlin DM III, Perez LR. Dietary effects on thyroid hormones in the red drum, *Sciaenops ocellatus*. *Fish Physiol Biochem* **11**:329–335, 1993.
137. Gorbman A, Dickhoff WW, Vigna SR, Clark NB, Ralph CL. *Comparative Endocrinology*. New York: Wiley, 1983.
138. Fletcher CR. The metabolism of iodine by a polychaete. *Comp Physiol Biochem* **35**:105–123, 1970.
139. Marcheggiano A, Iannoni C, Davoli C. Thyroglobulin-like activity in the nervous system of *Eisenia foetida* (Annelida, Oligochaeta). *Cell Tissue Res* **241**:429–433, 1985.
140. Falugi C, Marcheggiano A, Iannoni C, Davoli C. Localization of cholinesterase activity during the development of *Eisenia foetida*. *J Histochem Cytochem* **36**:949, 1988.
141. Davoli C, Marcheggiano A, Ravagnan G, Minu M, Serafino A, Iannoni C. Iodination activity in *Eisenia foetida* (Annelida, Oligochaeta). *Cell Tissue Res* **264**:9–14, 1991.
142. Anthéunisse LJ, Lever J. J131-accumulation in some invertebrates. *Koninklyke Nederlandse Akadamie Van Wetenschappen* **59**:562–565, 1956.
143. Major CW, Hanegan JL, Anoli L. Organic binding of iodide in nemertean mucus *in vivo* and *in vitro*. *Comp Biochem Physiol* **28**:1153–1160, 1969.
144. Roche J, Michel R. Natural and artificial iodoproteins. *Adv Prot Chem* **6**:253–297, 1951.
145. Roche J. Sur la biochimie comparée de l'iode et des hormones thyroïdiennes dans les organismes animaux. *Bull Soc Chim Biol* **42**:1155–1164, 1960.
146. Drechsel H. Beiträge zur Chemie einiger Seethiere. 11. Über das Achsenskelett der *Gorgonia cavellini*. *Z Biol* **33**:85–107, 1896.
147. Andre S. Sur la fixation et l'organification de l'iode par des hydres. *C R Soc Biol Paris* **159**:813–815, 1965.
148. Spangenberg DB. Iodine induction of metamorphosis in *Aurelia aurita*. *J Exp Zool* **165**:441–449, 1967.
149. Spangenberg DB. Thyroxine-induced strobilation in *Aurelia aurita*. *J Exp Zool* **178**:183–194, 1972.
150. Spangenberg DB. Thyroxine in early strobilation in *Aurelia aurita*. *Am Zool* **14**:825–831, 1974.
151. Spangenberg DB. Effects of exogenous thyroxine on statolith synthesis and resorption in *Aurelia*. *Am Zool* **24**:917–923, 1984.
152. Black RE, Webb KL. Metabolism of ^{131}I in relation to strobilation of *Chrysaora quinquecirrha* (Scyphozoa). *Comp Biochem Physiol A* **45**:1023–1029, 1973.
153. Olman S, Webb K. Metabolism of ^{131}I in relation to strobilation of *Aurelia aurita* L. (Scyphozoa). *J Exp Mar Biol Ecol* **16**:113–122, 1974.
154. Silverstone M, Tosteson TR, Cutress CF. The effect of iodide and various iodocompounds on initiation of strobilation in *Aurelia*. *Gen Comp Endocrinol* **32**:108–113, 1977.
155. Olman S, Webb K. Aspects of iodine metabolism in jellyfish polyps. *Va J Sci* **23**:114, 1972.
156. Silverstone M, Galton VA, Ingbar SH. Observations concerning the metabolism of iodine by polyps of *Aurelia aurita*. *Gen Comp Endocrinol* **34**:132–140, 1978.
157. Roche J, Yagi Y. Sur la présence de la monoiodotyrosine dans la spongine. *C R Soc Biol Paris* **146**:288–290, 1952.
158. Csaba G, Ubornyak L. Quantitative observations on triiodothyronine and histamine binding in *Tetrahymena*. *Acta Protozool* **18**:491–495, 1979.
159. Vought RL, Brown FA, Sibinovic KH, McDaniel EG. Effect of changing intestinal bacterial flora on thyroid function in the rat. *Horm Metab Res* **4**:43–47, 1972.
160. deHerder WW, Hazenberg MP, Pennock-Schroder AM, Oosterlaken AC, Rutgers M, Visser TJ. On the enterohepatic cycle of triiodothyronine in rats; Importance of the intestinal microflora. *Life Sci* **45**:849–856, 1989.
161. DiStefano JJ, DeLuze A, Nguyen T. Binding and degradation of 3,5,3'-triiodo-L-thyronine and thyroxine by rat intestinal bacteria. *Am J Physiol* **264**:E966–E972, 1993.
162. Hays MT. Thyroid hormones and the gut. *Endocr Res* **14**:203–224, 1988.
163. DiStefano JJ. Excretion, metabolism and enterohepatic circulation pathways and their role in overall thyroid hormone regulation in the rat. *Am Zool* **28**:373–387, 1988.

164. Salvatore G, Covelli I, Roche J. La fixation des hormones thyroïdiennes par *Escherichia coli* et son mécanisme. *Gen Comp Endocrinol* **3**:15–25, 1963.
165. Hisao SIC. Life history and iodine nutrition of the marine brown alga, *Petalonia fasciata* (O.F. Mull.) Kuntze. *Can J Bot* **47**:1611–1616, 1969.
166. Edmington J, Farmer M, Pitt-Rivers R, Ramsden DB, Rioch F. Thyroxine in plants. *Rec Adv Biol* **77**:131–138, 1994.
167. Tonkikh AK, Babaev TA, Markova IA, Gagelgans AI. Mechanism of auxin-like action of thyroxine. *Russian J Plant Physiol* **42**:170–175, 1995.
168. Martin CR. A Textbook of Endocrine Physiology. Baltimore: Williams and Wilkins, 1976.
169. Bell JM, Belzile RJ. Goitrogenic properties. In: Bowland GP, Clandinin DR, Wetter LR, Eds. Rapeseed Meal for Livestock and Poultry—A Review. Can Dept Agric Public No. 1257, Can. Dept. Agric., Ottawa. pp 45–60, 1966.
170. Higgs DA, McBride JR, Markert JR, Dosanjh BS, Plotnikoff MD, Clarke WC. Evaluation of Tower and Candle rapeseed (canola) meal and Bronowski rapeseed protein concentrate as protein supplements in practical dry diets for juvenile chinook salmon (*Oncorhynchus tshawytscha*). *Aquaculture* **29**:1–31, 1982.
171. Merken L, Simons MJ, Swillens S, Massaer M, Vassart G. Primary structure of bovine thyroglobulin deduced from the sequence of its 8,431-base pair complementary DNA. *Nature* **316**:647–651, 1985.
172. Mori N, Itoh N, Salvaterra PM. Evolutionary origin of cholinergic macromolecules and thyroglobulin. *Proc Natl Acad Sci U S A* **84**:2813–2817, 1987.
173. Parma J, Christophe D, Pohl V, Vassart G. Structural organization of the 5' region of the thyroglobulin gene. Evidence for intron loss and "exonization" during evolution. *J Mol Biol* **196**:769–779, 1987.
174. Swillens S, Ludgate M, Mercken L, Dumont JE, Vassart G. Analysis of sequence and structure homologies between thyroglobulin and acetylcholinesterase: Possible functional and clinical significance. *Biochem Biophys Res Commun* **137**:142–148, 1986.
175. Takagi Y, Omura T, Go M. Evolutionary origin of thyroglobulin by duplication of esterase gene. *Fed Eur Biochem Soc* **282**:17–22, 1991.
176. Varma SK, Collins M, Haller WS, Varma K. Thyroxine, triiodothyronine and reverse triiodothyronine concentrations in human milk. *J Pediatr* **93**:803–806, 1978.
177. Nizankowska-Blaz T, Abramowicz T, Korczowski R, Rusim J. Triiodothyronine and thyroxine in human, cow's and formula milk. *Exp Clin Endocrinol* **91**:116–118, 1988.
178. Gemmell RT, Sernia C. The transfer of thyroxine from the mother to the young of the marsupials, the bandicoot (*Isodon macrourus*) and the possum (*Trichosurus vulpecula*). *Comp Biochem Physiol A* **103**:541–543, 1992.
179. Morreale de Escobar G, Calvo R, Obregon MJ, Escobar del Rey F. Contribution of maternal thyroxine to fetal thyroxine pools in normal rats near term. *Endocrinology* **126**:2765–2767, 1990.
180. Tagawa M, Hirano T. Presence of thyroxine in eggs and changes in its content during early development of chum salmon, *Oncorhynchus keta*. *Gen Comp Endocrinol* **68**:129–135, 1987.
181. Prati M, Calvo R, Morreale de Escobar G. L-thyroxine and 3,5,3'-triiodothyronine concentrations in the chicken egg and in the embryo before and after the onset of thyroid function. *Endocrinology* **130**:2651–2659, 1992.
182. Specker JL. Preadaptive role of thyroid hormones in larval and juvenile salmon: Growth, the gut and evolutionary considerations. *Am Zool* **28**:337–349, 1988.
183. Evans ES, Schooley RA, Evans AB, Jenkins CA, Taurog A. Biological evidence for extrathyroidal thyroxine formation. *Endocrinology* **78**:983–1001, 1966.
184. Taurog A, Evans ES. Extrathyroidal thyroxine formation in completely thyroidectomized rats. *Endocrinology* **80**:915–925, 1967.
185. Obregon MJ, Mallol J, Escobar del Rey F, Morreale de Escobar G. Presence of L-thyroxine (T₄) and 3,5,3'-triiodo-L-thyronine (T₃) in tissues from thyroidectomized (T) rats. *Ann Endocrinol (Paris)* **42**:42, 1981.
186. Hedberg CW, Fishbein DB, Janssen RS, Meyers B, McMillen JM, MacDonald KL, White KE, Huss LJ, Hurwitz ES, Farhie JR, Simmons JL, Braverman LE, Ingbar SH, Schonberger LB, Osterholm MT. An outbreak of thyrotoxicosis caused by the consumption of bovine thyroid gland in ground beef. *N Engl J Med* **316**:993–998, 1987.
187. Bhatara VS, McMillin JM, Kummer M. Aggressive behavior and fire-setting in a four year-old boy associated with ingestion of ground beef contaminated with bovine thyroid tissue—A case report and review of neuropsychiatric findings in pediatric thyrotoxicosis. *J Child Adolesc Psychopharmacol* **5**:255–271, 1995.
188. Dymling J-F, Becker DV. Occurrence of hyperthyroidism in patients receiving thyroid hormone. *J Clin Endocrinol Met* **27**:1487–1491, 1967.
189. Chopra IJ. Nature, source and biological significance of thyroid hormones in blood. In: Werner SC, Ingbar SH, Eds. *The Thyroid*. Hagerstown, MD: Harper Row, pp 100–114, 1978.
190. Visser TJ, Mol JA, Otten MH. Rapid deiodination of triiodothyronine sulfate by rat liver microsomal fraction. *Endocrinology* **112**:1547–1549, 1983.
191. Matsuzaki S, Tonoue T, Yamamoto K. In vivo metabolism of iodotyrosines in thyroidectomized rats. *Endocrinol Jpn* **15**:64–69, 1968.
192. Maayan ML. Effect of growth hormone and thyrotropin upon deiodination of diiodotyrosine by hypophysectomized rat thyroid and liver. *Endocrinology* **78**:471–480, 1966.
193. Kobayashi I, Yamada T, Shichijo K. Effects of epinephrine and chemically related compounds on enzymatic deiodination of thyroxine, triiodothyronine, monoiodotyrosine and diiodotyrosine *in vitro*. *Metabolism* **15**:694–706, 1966.
194. Faberova A, Knopp J. Comparison of the *in vivo* deiodination of ¹²⁵I-, ¹²⁷I- and ¹³¹I-diiodotyrosine by thyroid and liver homogenates and slices from rat and pig. *Endocrinol Exp* **5**:245–251, 1971.
195. Eales JG. 1995 Measurement and regulation of thyroidal status in fish. *Neth J Zool* **45**:175–180, 1995.