

3647

Action of Narcotics on the Ameba by Means of Microinjection and Immersion.

STANISLAW HILLER.* (Introduced by Robert Chambers.)

From Cornell University Medical College, New York City.

Comparative studies were made on the influence of the narcotics ethyl alcohol, chloretone, ether, and chloroform on the protoplasm of *Amoeba dubia* with special reference to their action on the plasmalemma.

Immersion Experiments: Very weak concentrations (subnarcotic) cause the amebae to spread out and continue their movements in an expanded condition. This may be an effect of the lowered surface tension. Ether in its narcotic concentration (approximately 2.0 per cent) produces a reversible gelation accompanied by the cessation of all movements. Lethal concentrations of all the narcotics cause the ameba to round up, followed by a sinking of its granules and a disintegration of the plasmalemma.

Injection Experiments: No narcotic effect is produced by injection into the interior of the ameba. Chloretone in all concentrations increases the fluidity and streaming movements of the interior. Eighty per cent alcohol produces a coagulation which is localized and reversible when small amounts are injected and irreversible with large amounts. Chloroform and ether when injected into the cytoplasm form spherical drops which go into solution more rapidly than when injected into water. The protoplasm is irreversibly coagulated in the environment of the dissolving droplet. Chloroform is more potent than ether in causing coagulation. The irreversible coagula produced by the alcohol, chloroform and ether are pinched off by the living portion of the ameba.

Local Application of the Narcotic on the Exterior of the Ameba: With saturated chloretone solution and 95 per cent alcohol no local effects were obtained. All the pseudopodia are withdrawn and the ameba was converted into an actively moving limax form.

Concentrated, aqueous solution of ether produces a local blister on the plasmalemma with a coagulation of the cytoplasm at the base of the blister. A drop of pure ether dissolves the plasmalemma on remaining in contact with it for five seconds or more. This occurs most rapidly at the tip of a newly forming pseudopodium. As the ether droplet goes into solution there appears in its place a wrinkled

*Fellow International Education Board, Rockefeller Foundation.

pellicle enclosing several highly refractive globules. Both the pellicle and the globules are insoluble in water and in cold 95 per cent alcohol.

Chloroform similarly but more rapidly dissolves the plasmalemma. A small drop on touching the plasmalemma quickly passes through it and the endoplasm in its vicinity becomes coagulated.

An interesting phenomenon which possibly is related to the problem of pseudopodial formation occurs when chloroform is allowed to diffuse against the ameba from the tip of the micropipette. A local elevation of the pellicle takes place, into which the granular endoplasm flows. As the pipette is gradually withdrawn the incipient pseudopodium enlarges and extends in the direction of the moving pipette. In this way the entire ameba can be made to move to all appearances in a normal fashion along any path taken by the tip of the pipette.

I wish to thank Dr. Robert Chambers most sincerely for his kind help in this work.

3648

Influence of Vegetative Nerves Upon Distribution of Arsenic After Salvarsan Injections. II. Pilocarpinized Animals.

ERNST F. MÜLLER, C. N. MYERS AND E. MARPLES.

From the Department of Dermatology and Syphilology, College of Physicians and Surgeons, Columbia University.

A previous publication¹ gave a comparison between the distribution of arsenic after intravenous injection of arsenicals in normal animals² and in animals under the influence of epinephrin. When the involuntary nervous system is under the influence of epinephrin, almost 5 times as much arsenic is retained in the blood stream and only a small part of the arsenic normally distributed to the brain and liver reaches those organs. The general aspect of these tests led to the assumption that epinephrin inhibits the normal distribution of arsenic into the body organs.

In a second series of tests, 2 groups of rats were given subcutaneous injections of pilocarpin previous to the intravenous injections of salvarsan, neosalvarsan and silversalvarsan. 160 mg. of salvarsan per kg. body weight was the dosage used for the normal rats and the pilocarpinized rats were given 80 mg. per kg. body weight. The figures for neosalvarsan and silversalvarsan are similar to those