

Serum Cholinesterase Activity in Thymectomized Rats. (23083)

HUGH A. LINDSAY (Introduced by E. J. VanLiere)

Department of Physiology, School of Medicine, West Virginia University, Morgantown.

Hawkins *et al.*(1) found an inverse relation between thyroid and serum cholinesterase activities in the rat. Thyroidectomy produces regression of the thymus(2,3,4). Thyroid extract causes hypertrophy of the thymus, although an excess leads to atrophy(2). Then would thymectomy cause a change in esterase activity?

Methods. Fourteen young (50-60 g) female albino rats of the Wistar strain were thymectomized under light ether anesthesia according to the technic of Segaloff(5). Fourteen control rats were anesthetized and sham-operated; the incision was held open for the average time necessary to perform a thymectomy. Three weeks after the operations the animals were bled. Serum esterase activity was measured by a modification of the optical method developed by Kalow(6) (serum dilution, 1 to 100; temp. 40°C; substrate, 5×10^{-5} M benzoylcholine chloride; wavelength, 240 m μ ; duration of measurement, 15 min.).

Results were expressed as shift in absorbance $\times 1,000$.

Results. In the control rats the mean esterase activity was 37.5 units, in the thymectomized, 37.6 units.

Summary Although the thyroid gland influences both the status of the thymus and serum cholinesterase activity, thymectomy alone has no influence on serum cholinesterase activity.

1. Hawkins, R. B., Nichikawara, M., and Mendel, B., *Endocrinol.*, 1948, v43, 167.
2. Houssay, B. A., *Human Physiology*, McGraw-Hill, ed. 2, 1955.
3. Marine, D., Manley, O. T., and Baumann, E. J., *J. Exp. Med.*, 1924, v40, 429.
4. Chiodi, H., *Endocrinol.*, 1940, v26, 107.
5. Segaloff, A., cited by Farris, E. J., and Griffith, J. Q., Jr., *The Rat in Laboratory Investigation*, Lippincott, ed. 2, 1949.
6. Kalow, W., *J. Pharmacol. Exp. Therap.*, 1952, v26, 107.

Received February 11, 1957. P.S.E.B.M., 1957, v94.

Effect of Incubation upon Folic Acid Content of Eggs. (23084)

MIKE HAYES* AND THOMAS D. LUCKEY

Department of Biochemistry, School of Medicine, University of Missouri, Columbia.

The effect of incubation of eggs upon content of most of the B-vitamins has been determined. Snell and Quarles(1) found no change in the level of riboflavin or pantothenic acid in eggs throughout one period of incubation. Biotin concentration remained constant, or decreased slightly, while inositol and nicotinic acid were found to be synthesized by the embryo. Okamoto(2) found the concentration of thiamin to remain constant. Cielens(3) reports that 53-67% of choline is lost during

incubation. Folic acid has apparently not been studied.

Since germfree rats have been found to synthesize vit. B₉ (folic acid and citrovorum factor) while they require biotin and other B-vitamins(4), it is of special interest to determine the change in folic acid content of the incubating egg (which is usually germfree). Evans *et al.*(5) studied the effect of cold storage on folic acid content of eggs and Ebert(6) reported on development and role of p-aminobenzoic acid (a component of folic acid) in the early chick embryo.

Methods. Four dozen eggs were procured

* Medical Student Fellow of Nat. Fn. for Infantile Paralysis.