

***Brucella melitensis*, *B. neotomae* and *B. ovis* Elicit Common and Distinctive Macrophage Defense Transcriptional Responses**

JILL COVERT,*¹ ANGELA J. MATHISON,*¹ LINDA ESKRA,* MENACHEM BANAI,†
AND GARY SPLITTER*²

*Department of Pathobiological Sciences, University of Wisconsin-Madison, Madison, Wisconsin 53706; and †Kimron Veterinary Institute, Department of Bacteriology, Beit Dagan 50250, Israel

Brucella spp. establish an intracellular replicative niche in macrophages, while macrophages attempt to eliminate the bacteria by innate defense mechanisms. *Brucella* spp. possess similar genomes yet exhibit different macrophage infections. Few *B. melitensis* and *B. neotomae* enter macrophages with intracellular adaptation occurring over 4–8 hr. Conversely, *B. ovis* are readily ingested by macrophages and exhibit a persistent plateau of infection. Evaluating early macrophage interaction with *Brucella* spp. allows discovery of host entry and intracellular translocation mechanisms. Microarray analysis of macrophage transcriptional response following a 4 hr infection by different *Brucella* spp. revealed common macrophage genes altered in expression compared to uninfected macrophages. Macrophage infection with three different *Brucella* spp. provokes a common innate immune theme with increased transcript levels of chemokines and defense response genes and decreased transcript levels of GTPase signaling and cytoskeletal function that may affect trafficking of *Brucella* containing vesicles. For example, transcript levels of genes associated with chemotaxis (*IL-1β*, *MIP-1α*), cytokine regulation (*Socs3*) and defense (*Fas*, *Tnf*) were increased, while transcript levels of genes associated with vesicular trafficking (*Rab3d*) and lysosomal associated enzymes (*prosaposin*) were decreased. Genes with altered macrophage transcript levels among *Brucella* spp.

infections may correlate with species specific host defenses and intracellular survival strategies. Depending on the infecting *Brucella* species, gene ontology categorization identified genes differentially involved in cell growth and maintenance, endopeptidase inhibitor activity and G-protein mediated signaling. Examples of decreased gene expression in *B. melitensis* infection but not other *Brucella* spp. were growth arrest (*Gas2*), immunoglobulin receptor (*Fcγrl*) and chemokine receptor (*Cxcr4*) genes, suggesting opposing effects on intracellular functions. Exp Biol Med 234:1450–1467, 2009

Key words: *Brucella* infection; macrophages; transcriptome; *Brucella melitensis*; *Brucella ovis*; *Brucella neotomae*

Introduction

Brucella species (spp.) are zoonotic pathogens able to infect humans and cause abortion in domestic animals. Human infection generally requires contact with a limited number of organisms (infectious dose estimated as less than 100 organisms) (1), progresses with inconsistent and persistent flu-like symptoms from 2–6 weeks post-inoculation and, if left untreated, develops into chronic brucellosis. Macrophages phagocytose *Brucella* spp. and initiate an innate immune response, while *Brucella* subvert the host antimicrobial defense mechanisms to establish an intracellular replicative niche (2). Once resident within the macrophage, *Brucella* avoid exposure and killing by the humoral immune response.

Host preference and virulence among species are unaccounted when comparing the few differences between genomic sequences of *B. melitensis*, *B. abortus* and *B. suis* (3). When comparing six historically identified *Brucella* spp., only 217 open reading frames present in *B. melitensis* were absent in the other species (4). Ultimately, *Brucella* research has revealed a limited number of factors that significantly alter host specificity by *Brucella* spp. Human infections with *B. melitensis* are severe in pathogenesis and

This work was sponsored by the NIH/NIAID Regional Center of Excellence for Bio-defense and Emerging Infectious Diseases Research (RCE) Program. The authors wish to acknowledge membership within and support from the Region V Great Lakes RCE (NIH award 1-U54-AI-057153), R21 AI070229, and BARD US-3829–06.

¹ These authors contributed equally to the manuscript.

² To whom correspondence should be addressed at University of Wisconsin-Madison, 1656 Linden Dr., Madison, WI 53706. E-mail: Splitter@svm.vetmed.wisc.edu

Received April 1, 2009.
Accepted July 15, 2009.

DOI: 10.3181/0904-RM-124
1535-3702/09/23412-1450\$15.00
Copyright © 2009 by the Society for Experimental Biology and Medicine

are widely reported; conversely, neither *B. ovis* nor *B. neotomae* have been reported to cause human infection, and pathogenesis is at most limited. Investigating the murine macrophage response to highly similar *Brucella* spp. may provide additional understanding regarding the ability of *Brucella* spp. to establish and maintain infections.

Although transcriptional profiles of murine macrophages infected with *B. abortus* have been studied (7), no studies have compared host response among infections of differing *Brucella* spp. Altered host transcriptional response among *Brucella* spp. infections may identify not only common responses to infection, but also distinguish genes and pathways specific to each *Brucella* spp. infection. Identifying alterations in the macrophage transcriptome may provide greater understanding of host mechanisms involved in pathogen killing and bacterial regulation that limit damage to host cells during infection.

Murine macrophages are frequently used to investigate *Brucella* infection. The transcription profile after 4 hr of infection would evaluate general as well as specific response to different but genetically similar *Brucella* spp. Bacteria enter host cells and translocate to an endoplasmic reticulum containing a replicative niche within a few hours post infection; concurrently, a portion of the bacteria die by phagosome-lysosome fusion (8). The majority of host transcriptional response occurs during this early time (9). Examining an early time, such as 4 hr post infection, permits discovery of potential mechanisms of entry and intracellular translocation that take place before bacterial replication becomes evident after 8 hr (2, 10–12). The present microarray analyses evaluate macrophage response to *Brucella* spp. infection by testing 6 hypotheses and focusing on analogous and distinct transcriptional responses elicited by *B. melitensis*, *B. neotomae* and *B. ovis*.

Materials and Methods

Bacteria and Cell Lines. *B. melitensis*, *B. neotomae* and *B. ovis* were grown in 12- by 75-mm tubes on a shaker platform in BBL Brucella broth (BD Biosciences, Franklin Lakes, NJ) or on Brucella broth plates containing 1.5% agar. *B. melitensis*, *B. neotomae* and *B. ovis* were transformed with pBBR1MCS/GFP_{uv} containing green fluorescent protein (*gfp_{uv}*) under a constitutive Tac promoter and with chloramphenicol resistance (13). *Brucella* spp. for infections were grown in broth with or without chloramphenicol at 37°C for 1–2 days and colony forming units (CFUs) determined by plating on agar and incubating 3 days at 37°C with 5% CO₂.

RAW 264.7 (TIB-71, ATCC) and J774A.1 (TIB-67, ATCC) mouse macrophage cell lines were maintained at 37°C with 5% CO₂ in RPMI 1640 (Invitrogen, Carlsbad, CA) supplemented with 10% fetal bovine serum (FBS), 0.2 mM L-glutamine, antibiotic-antimycotic (100 U/mL penicillin G, 100 µg/mL streptomycin, 0.25 µg/mL amphotericin B, Gibco), 1 mM sodium pyruvate (SAFC Biosciences,

St. Louis, MO) and MEM amino acids (Hyclone, Logan, UT).

Intracellular Survival of *Brucella* spp. in Macrophages. Macrophages ($0.5\text{--}1 \times 10^6$ /well) were plated in 6-well plates 2–12 hr prior to infection in medium without antibiotics. *Brucella* spp. were grown to stationary phase in Brucella broth and then serially diluted and plated on Brucella agar to estimate CFU/mL. Macrophages were infected at a multiplicity of infection (MOI, bacteria to macrophage) of 1000:1 for 90 min at 37°C with 5% CO₂. Extracellular bacteria were removed using three PBS washes followed by 30 µg/mL gentamicin (MP Biomedicals, Inc., Irvine, CA) in RPMI. After 30 min, macrophages were washed three times with PBS. RPMI supplemented with 2 µg/mL gentamicin was added to cultures after 2 hr. At 4 hr, cultures were washed, lysed with 0.1% Triton X-100, serially diluted and plated twice on Brucella agar to determine bacterial CFUs. Experiments were repeated independently a minimum of three times.

Flow Cytometry of *Brucella*-GFP_{uv} Infected Macrophages. Macrophages (5×10^7 /T₂₅ flask) were plated 2–12 hr prior to infection in medium without antibiotics. *Brucella* spp. were grown to stationary phase, and macrophages were infected at a multiplicity of infection (MOI, bacteria to macrophage) of 1000:1 for 90 min at 37°C with 5% CO₂ followed by gentamicin treatment to remove extracellular bacteria as described above. After 4 hr of incubation, cells were washed three times with PBS, fixed in 4% paraformaldehyde (Electron Microscopy Sciences, Hatfield, PA) for 30 min and observed by fluorescence microscopy (Carl Zeiss, Thornwood, NY). Using a FACScan (Becton Dickinson, Palo Alto, CA), ten thousand events were collected. Debris and dead cells were eliminated from the analysis on the basis of side scatter and forward scatter. Cells infected with *B. melitensis*, *B. neotomae* and *B. ovis* strains containing pBBR1MCS/GFP_{uv} were analyzed by flow cytometry to determine percentage of RAW 264.7 or J774A.1 cells infected at 4 hr.

Macrophage Infection for RNA Isolation. RAW 264.7 cells for microarray and RAW 264.7 and J774A.1 cells for RT-PCR were plated at 5×10^6 cells/T₇₅ flask 12–24 hr prior to infection, in supplemented RPMI 1640 without antibiotics. Macrophages were infected with 1 mL of a stationary phase *Brucella* spp. culture (MOI 1000:1). Infected macrophages were incubated for 4 hr at 37°C with 5% CO₂, washed once with PBS and then lysed for RNA collection (RNeasy, Qiagen, Germantown, MD).

Target Preparation for Microarray Hybridization. Total RNA was isolated from macrophage cultures (RNeasy, Qiagen) with lysate centrifugation to remove intact bacterial cells. DNase treated RNA from two independent infections was pooled for each target preparation. Target RNA was prepared according to the manufacturer's protocols (Affymetrix, Santa Clara, CA). Briefly, RNA was converted to double stranded cDNA (Invitrogen) except that T-7-(dT)24 oligomer (Genset Corp., San Diego,

CA) was used. Double stranded cDNA was isolated using GeneChip® Sample cleanup and *in vitro* synthesis of biotin-labeled cRNA was completed with Enzo BioArray High-Yield RNA Transcript Labeling (Affymetrix). Labeled cRNA mixed with fragmentation buffer was incubated at 94°C for 35 min and was confirmed by agarose gel electrophoresis. Final RNA concentration ranged from 0.5–1.1 µg/µL.

Microarray Hybridization and Analysis. Labeled cRNA was hybridized to GeneChip® Test3 arrays and Murine Genome U74Av2 microarrays (Affymetrix). Eleven MG_U74Av2 GeneChip® microarrays were independently hybridized with cRNA from uninfected macrophage samples (2 independent samples) and each of three *Brucella* spp. infected macrophage samples (3 independent samples for each *Brucella* species infection). GeneChip® washing, hybridization and scanning was performed by the University of WI Biotechnology Center, Gene Expression Center (University of WI-Madison) utilizing Affymetrix protocols and procedures. Affymetrix *.CHP, *.CEL and spreadsheets of signal output are available through NCBI Gene Expression Omnibus database at the time of publication, Series accession number GSE8385.

mRNA from uninfected RAW 246.7 macrophages was compared to macrophages infected for 4 hr with *B. melitensis*, *B. ovis* or *B. neotomae*. All genes were subjected to analysis by EBarrays (14, 15), a statistical analysis package in the comprehensive R archive network (<http://cran.r-project.org/>). Data conformity to the statistical assumptions was checked (coefficient of variation and log-normal normal model) and best fit models were used. Six biologically relevant hypotheses of altered transcript levels were tested ($M\phi$ designates RAW 264.7 macrophages):

$$H_0: M\phi = M\phi + B. melitensis = M\phi + B. neotomae \\ = M\phi + B. ovis$$

Infected and uninfected macrophages have similar transcription.

$$H_1: M\phi \neq M\phi + B. melitensis = M\phi + B. neotomae \\ = M\phi + B. ovis$$

Brucella spp. infected macrophages express mRNA different from uninfected macrophages.

$$H_2: M\phi \neq M\phi + B. melitensis \neq M\phi + B. neotomae \\ = M\phi + B. ovis$$

B. ovis and *B. neotomae* infected macrophages express mRNA different from *B. melitensis* or uninfected macrophages.

$$H_3: M\phi \neq M\phi + B. melitensis = M\phi + B. neotomae \\ \neq M\phi + B. ovis$$

B. melitensis and *B. neotomae* infected macrophages

express mRNA different from *B. ovis* or uninfected macrophages.

$$H_4: M\phi \neq M\phi + B. melitensis = M\phi + B. ovis \\ \neq M\phi + B. neotomae$$

B. melitensis and *B. ovis* infected macrophages express mRNA different from *B. neotomae* or uninfected macrophages.

$$H_5: M\phi \neq M\phi + B. melitensis \neq M\phi + B. neotomae \\ \neq M\phi + B. ovis$$

Each macrophage culture expresses distinct mRNA.

Empirical Bayesian statistics designated a posterior probability (EBarrays Probability) specific to each gene, evidence that a given gene follows the transcription pattern designated by each of the hypotheses. The probability of random gene assignment to any hypothesis was 0.167, while posterior probabilities ≥ 0.2 were considered significant. Genes were sorted within each experimental hypothesis, and each gene was allocated only to the hypothesis with the largest posterior probability. The signal logarithm ratio (SLR) was calculated as the logarithm, base 2, of the ratio between infected (experimental) signal and uninfected (control) signal. Thus, the experimental:control ratio was transformed to zero for no change between conditions and the equivalent of a two-fold increase or decrease becomes 1 or -1 SLR, respectively.

Additional data analysis was conducted utilizing TM4 microarray analysis tools ANOVA (16) and SAM (17, 18) contained within TIGR MultiExperiment Viewer (MeV) (<http://www.tm4.org/>) (19, 20). Genes were annotated utilizing Affymetrix's NetAffx Analysis Center (<http://www.affymetrix.com/analysis/index.affx>) and DAVID Bioinformatics Resources (<http://david.abcc.ncifcrf.gov/>). Gene ontology (GO) categorization of genes identified with altered transcription between uninfected and *Brucella* spp. infected macrophages were completed with EASE (<http://david.abcc.ncifcrf.gov/>). Genes were iteratively subjected to GO biological processes categorization and, while genes may be categorized in several of the GO categories, each gene was listed once.

Real Time RT-PCR (RT-PCR). Total RNA was isolated from macrophage cultures (RNeasy, Qiagen) according to manufacturer's protocol with DNase treatment. Macrophage RNA (2–4 µg) was primed with oligo(dT)₂₀, reverse transcribed with SuperScript™ II or III (Invitrogen), and resultant cDNA diluted 1:5. Primers (listed in Text S1) and double-dye oligonucleotide probes for PCR were developed to amplify mRNA fragments 80–130 basepairs in length using Primer3 (21). RT-PCR was performed on cDNA samples utilizing qPCR Mastermix (Eurogentec, San Diego, CA) or iQ supermix (Bio-Rad, Hercules, CA) with dual labeled probes and iQ SYBR® green Supermix (Bio-Rad), respectively, to quantify




Figure 1. Intracellular survival of *Brucella* spp. in RAW 264.7 and J774A.1 macrophages. Panel A. Intracellular *Brucella* spp. were isolated from RAW 264.7 or J774A.1 macrophages and enumerated at 4 hr. Macrophages were infected with 1000:1 MOI of each *Brucella* spp. and allowed to infect for 90 min followed by gentamicin treatment for 30 min. At 4 hr, macrophages were washed then lysed with 0.1% Triton X-100. Intracellular *Brucella* were enumerated by plating serial dilutions on agar. Four independent experiments were conducted, and the number of *Brucella* spp. isolated per well was averaged with error bars representing the SEM. Comparing *B. melitensis* or *B. neotomae* to *B. ovis* infection is significantly different with $P \leq 0.005$. Panel B. Flow cytometry analysis of *Brucella*-GFP_{uv} infection in RAW 264.7 and J774A.1 macrophages at 4 hr. RAW 264.7 or J774A.1 macrophages were infected for 90 min with *Brucella* spp. containing GFP_{uv} followed by gentamicin treatment to remove extracellular bacteria. After 4 hr, RAW 264.7 or J774A.1 cells were fixed in 4% paraformaldehyde, and the percent of infected cells determined by flow cytometry analysis. Comparing *B. melitensis* or *B. neotomae* to *B. ovis* infection is significantly different with $P \leq 0.005$. Panel C. RAW 264.7 macrophages were infected continuously for 4 hr with *Brucella* spp. containing GFP_{uv}. Cells were washed to remove extracellular *Brucella* and fixed in 4% paraformaldehyde. Matched bright field and fluorescence images were digitally captured at ×63 oil immersion magnification. Bar equals 20 μ m. A color version of this figure is available in the online journal.

relative transcript levels. The PCR was optimized on the iCycler (Bio-Rad) with primer concentrations ranging from 300–500 nM and probe concentrations at 125 nM. Amplification cycles were 95°C for 15 sec followed by 60 sec at 60°C with fluorescence detected during the extension phase. Relative gene mRNA quantities were quantified by the standard curve method using a house-keeping gene as a reference gene. SLR was calculated with these normalized transcription values.

ELISA Measurement of TNF Activity. BMDM cells were flushed from the femurs and tibiae of 10–12 week-old wild-type C57BL/6 mice. Cells were grown for 5–8 days in RPMI 1640 (Gibco, Grand Island, NY), 10% FBS (Equitech-Bio Inc., Kerrville, TX), 25 μ g/mL gentamicin

(MP Biomedicals, Inc., Solon, OH) and 20–30 ng/mL M-CSF (R&D Systems, Minneapolis, MN). BMDM cells plated at 1×10^6 cells/well in 6-well plates were infected with *B. melitensis* at an MOI 100 and incubated at 37°C. Supernatants were collected at 12 hr from cells infected with *B. melitensis* or medium. Supernatants were filtered through a 0.22 micron PES Millipore filter (Millipore, Billerica, MA) and assayed for TNF α using a Ready-Set-Go ELISA kit (eBioscience, San Diego, CA).

Additional Calculations and Statistics. Error bars throughout indicate the standard error of the mean (SEM). A Student's *t* test determined if there was significant difference between two independent samples using a pooled estimate of variation.

Table 1. RAW 264.7 Macrophage Transcription of Common Housekeeping Genes Following *Brucella* spp. Infection

H ₀ : M ϕ = M ϕ + <i>B. melitensis</i> = M ϕ + <i>B. neotomae</i> = M ϕ + <i>B. ovis</i>						
Probe set	Entrez gene	Average signal	SLR ^a	EBarrays probability	Gene symbol	Gene product
101213_at	11837	5.7×10^4	-0.2 ± 0.06	1.000	Arbp	Acidic ribosomal phosphoprotein PO
101578_f_at	11461	2.2×10^4	-0.6 ± 0.6	0.870	Actb	β -actin
101214_f_at 162210_r_at 97751_f_at	14433	2.5×10^4	0.1 ± 0.2	0.894	Gapdh	Glyceraldehyde-3-phosphate dehydrogenase
93346_at	18655	2.0×10^4	0.2 ± 0.1	1.000	Pgk1	Phosphoglycerate kinase 1
93088_at	12010	1.6×10^4	0.2 ± 0.2	1.000	b2m	β 2 microglobulin
97538_at	110006	6.8×10^3	-0.6 ± 0.2	0.999	b-gus	β glucuronidase
160107_at	15452	6.5×10^3	0.07 ± 0.1	0.999	Hprt1	Hypoxanthine guanine phosphoribosyl transferase 1
103957_at 103958_g_at	22042	3.4×10^3	-0.5 ± 1.2	0.991	Tfrc	Transferrin receptor
99950_at	21374	6.7×10^2	-0.09 ± 0.2	0.996	Tbp	TATA box binding protein

^a Signal log ratio = LOG₂ (experimental *Brucella* spp. infected signal/control uninfected signal) $\pm$ SE.

Results

***Brucella* spp. Infection of RAW 264.7 and J774A.1 Macrophages Assessed by Colony Forming Units (CFUs) and Flow Cytometry Analysis of *Brucella*-GFP_{uv}.** RAW 264.7 macrophage infections with *B. melitensis* and *B. neotomae* were similar, with low numbers of these bacteria phagocytosed compared to the number of bacteria introduced (Fig. 1, panel A). Macrophage uptake of *B. ovis*, at 4 hr in both RAW 264.7 and J774A.1 cells was nearly two logarithmic units higher than *B. melitensis* or *B. neotomae*, paralleling previous observations (6, 22).

Similar to CFU results, flow cytometry analysis of RAW 264.7 macrophages infected for 4 hr with *Brucella* spp. expressing GFP_{uv} had low levels of *B. melitensis* and *B. neotomae*, compared to *B. ovis* (Fig. 1, panel B). This observation was further supported by fluorescent microscopy (Fig. 1, panel C). At this early time, 20–30 percent of macrophages were infected with *B. melitensis* or *B. neotomae*, while greater than 95 percent of macrophages were infected with *B. ovis* (Fig. 1, panel B) consistent with uptake for smooth and rough *Brucella* (22). Interestingly, the numbers of *Brucella* phagocytosed at these levels did not correlate in a linear or proportional manner to levels of transcriptional response, as changes in macrophage mRNA amounts following *B. ovis* infection were often similar to *B. melitensis* and *B. neotomae* infections.

Because of the difference in macrophage infection by *B. melitensis* and *B. neotomae* compared to *B. ovis*, macrophage viability was examined to determine if infection levels would influence macrophage survival. RAW 264.7 and J774A.1 macrophages were continuously infected with *Brucella* spp. for 4 hr without antibiotics and viability determined. No significant macrophage death was observed (data not shown). Therefore, host transcription with *Brucella* spp. infection at 4 hr was not biased toward cell death pathways.

Microarray Analysis of RAW 264.7 Macrophages Infected with Each of Three *Brucella* spp. *Genes Identified for Housekeeping Controls with No Transcriptome Change Across Three *Brucella* spp. Infection Conditions.* A series of commonly used housekeeping genes were evaluated for potential as control genes in downstream analysis, and changes in transcript levels between *Brucella* spp. infected and uninfected macrophages were compared (Table 1). Genes with high posterior probabilities and low signal log ratio (SLR) with small variability appeared as the best control genes for *Brucella* spp. infections. *Hypoxanthine guanine phosphoribosyl transferase 1* (*Hprt1*) and the *TATA box binding protein* (*Tbp*), which function in glycolysis and transcription, respectively (23), were transcribed at a moderate level (signal strength on microarray between 1×10^4 and 1×10^2) and had SLR near zero with small variability. *Hprt1* and *Tbp* were unaltered by *Brucella* spp. infections and were therefore utilized as control housekeeping genes in downstream analysis of the transcriptome. The transcript levels of β -actin, β -glucuronidase, *transferrin receptor* and *glyceraldehyde-3-phosphate dehydrogenase* were minimally altered (-0.6 to 0.1 SLR) by infection, but each had a pronounced standard error and were considered not optimal for use as control genes.

*Genes Identified With Increased Transcription Following *Brucella* spp. Infections.* Seventy-two genes were identified under H₁ (transcription from macrophages infected by three *Brucella* spp. differs from uninfected macrophages) with SLR increases between 2.0 and 6.8 following infection with any *Brucella* spp. (Table 2). Defense and chemotactic response, both related to a common inflammatory response, encompass the largest groups of genes with increased transcript levels during infection. *Interleukin-1 β* , *tumor necrosis factor*, macrophage inflammatory protein genes (*MIP-1 α* , *MIP-1 β* , *MIP-2 α*), colony stimulating factor genes (*Csf2* and *Csf3*) and *Fas* had increased transcription similar to reports with a

Table 2. Genes with Increased Transcription in *Brucella* spp. Infected as Compared to Uninfected RAW 264.7 Macrophages

H ₁ : MΦ ≠ MΦ + <i>B. melitensis</i> = MΦ + <i>B. neotomae</i> = MΦ + <i>B. ovis</i>									
Probe set	Entrez gene	SLR ^a			EBarrays probability	Gene symbol	Gene product	Additional statistics ^c	
		M ^b	N ^b	O ^b					
Chemotaxis									
102736_at	20296	6.4	5.8	6.5	0.996	Ccl2	Chemokine (C-C motif) ligand 2 (MCP-1)	a, b, c	
103486_at	16176	7.5	6.1	6.7	0.869	Il1b	Interleukin 1 beta	a, b, c	
94761_at	20306	5.6	4.6	6.0	0.764	Ccl7	Chemokine (C-C motif) ligand 7 (MCP-3)	a, b, c	
98406_at	20304	2.7	2.2	3.8	0.658	Ccl5	Chemokine (C-C motif) ligand 5 (RANTES)		
101160_at	20310	4.8	4.4	4.6	0.997	Cxcl2	Chemokine (C-X-C motif) ligand 2 (MIP-2α)	a, b, c	
941146_at	20303	3.6	3.3	3.6	0.999	Ccl4	Chemokine (C-C motif) ligand 4 (MIP-1β)	a, b, c	
98822_at	53606	2.3	2.0	3.0	0.891	lsg15	ISG15 ubiquitin-like modifier		
102424_at	20302	2.6	2.4	2.9	0.905	Ccl3	Chemokine (C-C motif) ligand 3 (MIP-1α)	a, b	
104388_at	20308	2.8	2.4	2.7	0.932	Ccl9	Chemokine (C-C motif) ligand 9 (MIP-1γ)	a, b	
93858_at	15945	1.7	2.1	2.2	0.522	Cxcl10	Chemokine (C-X-C motif) ligand 10	c	
Defense response									
92948_at	12981	6.4	5.0	5.1	0.723	Csf2	Colony stimulating factor 2 (granulocyte-macrophage)	a, b	
100981_at	15957	4.6	4.2	5.6	0.720	Ifit1	Interferon-induced protein with tetratricopeptide repeats 1	a, b, c	
102629_at	21926	4.3	4.1	4.2	0.999	Tnf	Tumor necrosis factor	a, b, c	
94142_at	12985	4.7	3.7	4.6	0.987	Csf3	Colony stimulating factor 3 (granulocyte)	a, b, c	
103639_at	15958	2.8	3.2	3.7	0.907	Ifit2	Interferon-induced protein with tetratricopeptide repeats 2		
93871_at	16181	4.0	3.4	3.9	0.996	Il1m	Interleukin 1 receptor antagonist	a, b, c	
102921_s_at	14102	2.5	3.5	2.7	0.914	Fas	Fas (TNF receptor superfamily member)	c	
102712_at	20210	3.4	2.3	2.9	0.948	Saa3	Serum amyloid A 3	c	
94928_at	21938	2.5	2.0	2.5	0.885	Tnfrsf1b	Tumor necrosis factor receptor superfamily, member 1b	a, b, c	
92534_at	14579	2.6	2.0	2.9	0.839	Gem	GTP binding protein (gene overexpressed in skeletal muscle)		
98988_at	80859	3.1	2.7	2.1	0.94304859	Nfkbi2	Nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, zeta		
Protein-nucleus import									
97238_at	21335	2.8	3.2	2.9	0.940	Tacc3	Transforming, acidic coiled-coil containing protein 3		
104149_at	18035	2.6	2.1	1.9	0.704	Nfkbia	Nuclear factor of kappa light chain gene enhancer in B-cells inhibitor, alpha	a, b	
161135_f_at	66406	2.3	1.2	2.6	0.43736716	Sac3d1	SAC3 domain containing 1		
Apoptosis									
94186_at	22029	5.0	4.4	4.6	0.943	Traf1	Tnf receptor-associated factor 1	a, b, c	
104712_at	17869	3.7	4.0	3.0	0.890	Myc	Myelocytomatosis oncogene	c	
99392_at	21929	2.6	2.3	2.0	0.814	Tnfrsf3	Tumor necrosis factor, alpha-induced protein 3	c	
161666_f_at	17873	3.8	3.4	3.6	0.970	Gadd45b	Growth arrest and DNA-damage-inducible 45 beta	a, b	
102779_at	17873	2.7	2.4	2.7	0.947	Gadd45b	Growth arrest and DNA-damage-inducible 45 beta	c	
Fatty acid biosynthesis									
104647_at	19225	5.6	4.7	4.4	0.929	Ptgs2	Prostaglandin-endoperoxide synthase 2	a, b, c	
94057_g_at	20249	2.2	2.4	2.2	0.849	Scd1	Stearyl-coenzyme A desaturase 1		

Table 2. (Continued)

H ₁ : Mφ ≠ Mφ + <i>B. melitensis</i> = Mφ + <i>B. neotomae</i> = Mφ + <i>B. ovis</i>									
Probe set	Entrez gene	SLR ^a				EBarrays probability	Gene symbol	Gene product	Additional statistics ^c
		M ^b	N ^b	O ^b	Average ± SEM				
Cell surface receptor linked signal transduction									
95344_at	16165	3.0	3.3	3.0	3.1 ± 0.3	0.941	Il13ra2	Interleukin 13 receptor, alpha 2	c
97733_at	11541	2.8	2.6	2.5	2.7 ± 0.2	0.931	Adora2b	Adenosine A2b receptor	a, b
104498_at	26556	2.4	3.2	1.3	2.3 ± 0.3	0.421	Homer1	Homer homolog 1 (<i>Drosophila</i>)	b
102663_at	18793	2.7	2.2	2.7	2.5 ± 0.2	0.953	Plaur	Urokinase plasminogen activator receptor	a, b, c
Regulation of biological process									
92232_at	12702	5.6	4.4	5.2	5.1 ± 0.2	0.912	Socs3	Suppressor of cytokine signaling 3	a, b, c
162206_f_at	12702	3.9	3.1	3.4	3.5 ± 0.1	0.967	Socs3	Suppressor of cytokine signaling 3	a, b
92471_i_at	20556	2.4	1.6	2.6	2.2 ± 0.3	0.595	Slfn2	Schlafen 2	
Regulation of transcription, DNA-dependent									
101415_i_at	81845	2.5	2.9	1.9	2.4 ± 0.3	0.777	Bat4	HLA-B associated transcript 4	
102709_at	15260	2.4	2.5	2.7	2.5 ± 0.1	0.915	Hira	Histone cell cycle regulation defective homolog A (<i>S. cerevisiae</i>)	a, b
103651_r_at	68705	2.4	2.2	2.0	2.2 ± 0.2	0.808	Gtf2f2	General transcription factor IIF, polypeptide 2	
102882_at	22704	2.4	2.1	2.5	2.3 ± 0.3	0.579	Zfp46	Zinc finger protein 46	
Protein metabolism									
160829_at	21664	5.1	4.7	4.5	4.8 ± 0.4	0.958	Phlda1	Pleckstrin homology-like domain, family A, member 1	a, b
93352_at	21817	2.2	1.8	2.0	2.0 ± 0.5	0.530	Tgm2	Transglutaminase 2, C polypeptide	
102782_at	71340	3.3	3.5	2.8	3.2 ± 0.2	0.926	Rio1	RIO kinase 1	a, b
97548_at	328110	3.1	2.7	2.9	2.9 ± 0.2	0.950	Prpf39	PRP39 pre-mRNA processing factor 39 homolog (yeast)	
Cell growth and/or maintenance									
104451_at	18174	2.4	1.8	2.4	2.2 ± 0.1	0.805	Slc11a2	Solute carrier family 11 (proton-coupled divalent metal ion transporters), member 2 (Nramp2)	a, b
94379_at	16561	3.2	2.8	3.2	3.1 ± 0.1	0.945	Kif1b	Kinesin family member 1B	a, b
94384_at	15937	2.8	2.6	2.2	2.5 ± 0.3	0.890	Ier3	Immediate early response 3	b, c
160729_f_at	21884	3.3	2.1	3.4	2.9 ± 0.2	0.736	Fabp9	Fatty acid binding protein 9, testis	a, b
161281_f_at	15937	3.6	3.0	1.8	2.8 ± 0.3	0.44599685	—	Immediate early response 3	
Physiological process									
160084_at	546355	3.2	2.7	3.1	3.0 ± 0.3	0.995	Odc	Similar to ornithine decarboxylase	b
AFFX-GapdhMur/M	14433	2.9	2.3	1.7	2.3 ± 0.4	0.761	Gapdh	Similar to glyceraldehyde-3-phosphate dehydrogenase	
32599_5_st									
102749_at	12865	1.9	2.4	2.5	2.3 ± 0.4	0.672	Cox7a1	Cytochrome c oxidase, subunit VIIa 1	
94147_at	18787	2.6	2.3	1.9	2.3 ± 0.3	0.813	Serpine1	Serine (or cysteine) proteinase inhibitor, clade E, member 1	c
102694_at	26436	2.5	1.5	3.4	2.5 ± 0.2	0.498	Psg16	Pregnancy specific glycoprotein 16	a, b

Table 2. (Continued)

$H_1: M\phi \neq M\phi + B. melitensis = M\phi + B. neotomae = M\phi + B. ovis$									
Probe set	Entrez gene	SLR ^a			EBarrays probability	Gene symbol	Gene product	Additional statistics ^c	
		M ^b	N ^b	O ^b					
Miscellaneous classification									
96515_at	14204	1.9	1.9	2.5	2.1 ± 0.4	Il4i1	Interleukin 4 induced 1	a, b, c a, b	
98774_at	16365	4.0	4.2	4.1	4.1 ± 0.3	Irg1	Immunoresponsive gene 1		
98773_s_at	16365	3.4	3.1	3.3	3.3 ± 0.3	Irg1	Immunoresponsive gene 1		
94971_at	72391	2.7	1.9	2.6	2.4 ± 0.4	Cdkn3	Cyclin-dependent kinase inhibitor 3		
162384_f_at	12457	1.9	3.4	2.8	2.7 ± 0.4	Ccrn4l	CCR4 carbon catabolite repression 4-like		
94389_at	66373	1.9	2.4	1.9	2.1 ± 0.4	Lsm5	LSM5 homolog, U6 small nuclear RNA associated (S. cerevisiae)	b b b, c a, b	
97714_r_at	54130	2.7	1.7	3.2	2.5 ± 0.3	Actr1a	ARP1 actin-related protein 1 homolog A (yeast)		
96162_at	50764	1.6	3.4	2.3	2.5 ± 0.2	Fbxo15	F-box only protein 15		
93869_s_at	12044	2.3	2.2	2.3	2.3 ± 0.2	Bcl2a1a	B-cell leukemia/lymphoma 2 related protein A1a		
94505_at	67245	1.9	1.8	2.2	2.0 ± 0.1	Pel1	Pellino 1		
104177_at	58185	2.8	1.9	3.0	2.5 ± 0.3	Rsad2	Radical S-adenosyl methionine domain containing 2	b b b, c a, b	
100669_at	25465	2.1	2.0	2.5	2.2 ± 0.6	—	Interleukin 17		
161511_f_at	53606	2.1	2.1	3.1	2.4 ± 0.5	Isg15	Interferon stimulated gene, ubiquitin-like modifier		
97693_at	30865	5.6	5.0	5.4	5.3 ± 0.2	C78513	EST C78513		
104477_at	—	3.0	3.3	3.8	3.4 ± 0.3	—	Transcribed locus		
99849_at	319202	2.7	2.0	2.7	2.5 ± 0.2	1200016	RIKEN cDNA 1200016E24 gene	E24Rik	
					0.908				

^a Signal log ratio = LOG_2 (experimental *Brucella* spp. infected signal/control uninfected signal) $\pm$ SEM.^b *M*, *N*, *O* indicate *Brucella* spp. used to infect macrophages *B. melitensis*, *B. neotomae* and *B. ovis*, respectively.^c Identified by additional statistical analyses: *a*, ANOVA $P \leq 0.01$; *b*, SAM $\Delta = 0.05$; *c*, *B. abortus* microarrays (14).

variety of infectious agents (7, 24). Also, increased transcription was observed for cytokine regulation, anti-inflammatory and/or anti-apoptotic response genes (*Socs3*, *Slfn2*, *IL-1rn*, *Gadd45b* and *Tnfaip3*). These genes may prevent commitment to an inflammatory response pathway camouflaging immune recognition and providing a safe environment for bacterial survival. Although macrophages were infected with a greater number of *B. ovis* compared to *B. melitensis* and *B. neotomae*, Figure 1, no statistical difference was observed in the SLR between the 72 macrophage genes of Table 2 when infected with *B. ovis* compared to *B. melitensis* or *B. neotomae*.

Genes Identified with Decreased Transcription Following *Brucella* spp. Infections. Fifty-eight genes were identified with SLR decreases between -2.0 and -3.4 following infection with any *Brucella* spp. (Table 3). GO categories for genes with repressed transcript levels represented several biological functions, ranging from small GTPase mediated signal transduction and carboxylic acid metabolism to cell proliferation and lysosomal proteins. Small GTPase mediated signaling may be altered in response to the engulfment and association of *Brucella* spp. with the membrane of macrophages; included among this group are *Rab3d*, *Gna12*, *Cfl2* and *Iqgap1*. Small GTPases are key regulators associated with trafficking of *Brucella* containing vesicles to the endoplasmic reticulum (25), a crucial step in the establishment of a replication niche. In contrast to the inflammatory response genes commonly increased following infection, there were also genes categorized as response to external stimulus with decreased transcription. *Mr1*, *Abhd8* and *IL17a* may provide insight into mechanisms that the macrophage does not use in the response against this intracellular bacterium. Also, a decrease in *prosaposin* transcript levels was observed, a lysosomal enzyme that catabolizes glycosphingolipids that may enhance intracellular survival of *Brucella* spp.

Genes Identified with Altered Transcription Among *Brucella* spp. Infections. Thirty-three genes (Table 4) were identified under hypotheses that transcription among *Brucella* spp. infections was different from each other and uninfected macrophages (H_{2-4}). Of the 33 genes with altered expression among *Brucella* spp. infections, 21 genes had decreased transcript levels, 10 genes had increased transcript levels and, for two genes, the direction of change in expression was dependent on the species used for infection. Generally, the direction of change in transcription was uniform across the species tested. Twelve genes were altered based on the virulence descriptions of *Brucella* spp. (Table 4, part A), as *B. melitensis* was compared to *B. neotomae* and *B. ovis* infections (H_2). GO biological and molecular function categorization unveiled genes altered according to H_2 are involved in cell growth and maintenance, endopeptidase inhibitor activity, response to external stimuli and G-protein mediated signaling. When the infections were grouped according to LPS phenotype of the *Brucella*, i.e., smooth *B. melitensis* and *B. neotomae*

versus rough *B. ovis*, eleven genes were identified with altered transcription (H_3) (Table 4, part B). Although smooth *Brucella* enter macrophages via lipid rafts (26), the mechanism of *B. ovis* entry is unknown. If different entry mechanisms occur for *B. ovis*, a distinct group of host genes may be altered compared to smooth *Brucella* spp. However, few macrophage genes were altered between rough and smooth *Brucella* spp. infections, and there was little GO commonality. Ten genes had altered transcript levels when infection with *B. melitensis* and *B. ovis* were considered similar and distinct from *B. neotomae* infection (H_4) (Table 4, part C). All three *Brucella* spp. infections increased *interleukin 1 alpha* gene expression, but levels were higher in *B. melitensis* and *B. ovis* infections. Genes identified according to this pattern (H_4) were categorized in GO biological functions of cell growth and maintenance and signal transduction. Lastly, no genes were identified with each of the four conditions, uninfected and three *Brucella* spp. infections, having distinct transcription (H_5).

Confirmation of Microarray Data by Quantitative Real Time-PCR (qRT-PCR). To verify the microarray changes in transcript levels, eight genes were analyzed by qRT-PCR using the infection conditions as for microarray analysis. Housekeeping genes, *Hprt* and *Tbp*, identified via this microarray analysis and previously tested (23, 27, 28) were used to normalize cDNA levels and resulted in similar patterns and levels of gene transcription (data not shown). qRT-PCR confirmed that the eight genes tested (*Socs3*, *Tnf*, *IL-1 α* , *IL-1 β* , *Fc γ RI*, *Mef2c*, *Cxcr4* and *Abcd2*) had altered expression in RAW 264.7 macrophages following *Brucella* spp. infection in the same direction (increased or decreased) indicated by microarray analysis (Fig. 2, panels A and B). Further, J774A.1 and RAW 264.7 macrophage cell lines were used in parallel to verify that qRT-PCR results found in RAW 264.7 were applicable to J774A.1 cells. RAW 264.7 gene transcription altered among *Brucella* spp. infections was similarly changed (increased or decreased) during J774A.1 infection (Fig. 2, panels C and D) with the direction of altered transcription paralleling microarray results.

TNF α Production by Macrophages. To confirm the microarray change in TNF α transcription, bone marrow derived macrophages were infected with *B. melitensis* and supernatant analyzed by ELISA. Figure 3 illustrates that infected macrophages produced greater than 15-fold more TNF α than non-infected macrophages. These results together with the microarray findings suggest that *B. melitensis* activates TNF α production by macrophages.

Discussion

Murine macrophages infected with *Brucella* strains were profiled by microarray analysis. Interestingly, *B. melitensis* causes severe human infection, but neither *B. ovis* nor *B. neotomae* cause human pathogenesis. However, *in vitro* growth of *B. melitensis* and *B. neotomae* is similar in

Table 3. Genes with Decreased Level of Transcript in *Brucella* spp. Infected as Compared to Uninfected RAW 264.7 Macrophages

H ₁ : M ϕ $\neq$ M ϕ + <i>B. melitensis</i> = M ϕ + <i>B. neotomae</i> = M ϕ + <i>B. ovis</i>									
Probe set	Entrez gene	SLR ^a			Average $\pm$ SEM	EBarrays probability	Gene symbol	Gene product	Additional statistics ^c
		M ^b	N ^b	O ^b					
Small GTPase mediated signal transduction									
97415_at	19340	-2.6	-2.0	-3.2	-2.6 $\pm$ 0.3	0.866	Rab3d	RAB3D, member RAS oncogene family	b, c
97227_at	14673	-2.2	-2.5	-1.9	-2.2 $\pm$ 0.3	0.807	Gna12	Guanine nucleotide binding protein, alpha 12	a
97549_at	12632	-2.1	-2.2	-1.9	-2.1 $\pm$ 0.3	0.662	Cfl2	Cofilin 2, muscle	
93850_at	29875	-2.7	-2.0	-1.3	-2.0 $\pm$ 0.4	0.494	Iqgap1	IQ motif containing GTPase activating protein 1	
Cell proliferation									
161417_r_at	18109	-2.5	-2.8	-2.5	-2.6 $\pm$ 0.5	0.968	Mycn	V-myc myelocytomatosis viral related oncogene, neuroblastoma derived (avian)	
93943_f_at	12193	-2.5	-2.4	-2.4	-2.4 $\pm$ 0.5	0.925	Zfp36l2	Zinc finger protein 36, C3H type-like 2	
93713_at	20416	-2.5	-2.7	-2.2	-2.5 $\pm$ 0.3	0.914	Shc1	Src homology 2 domain-containing transforming protein C1	
92300_at	17428	-2.4	-2.0	-2.9	-2.4 $\pm$ 0.3	0.889	Mnt	Max binding protein	c
99076_at	353187	-2.2	-1.7	-2.8	-2.3 $\pm$ 0.3	0.786	Nr1d2	Nuclear receptor subfamily 1 group D member 2	c
101571_g_at	16010	-2.0	-1.7	-3.0	-2.3 $\pm$ 0.2	0.685	Igfbp4	Insulin-like growth factor binding protein 4	
99024_at	17122	-2.0	-1.8	-2.2	-2.0 $\pm$ 0.3	0.557	Mad4	Max dimerization protein 4	
100444_at	12568	-2.4	-1.9	-1.6	-2.0 $\pm$ 0.4	0.520	Cdk5	Cyclin-dependent kinase 5	
Carboxylic acid metabolism									
93320_at	12894	-3.4	-1.9	-3.1	-2.8 $\pm$ 0.4	0.663	Cpt1a	Camitine palmitoyltransferase 1a, liver	c
96799_at	30839	-2.7	-2.2	-2.3	-2.4 $\pm$ 0.3	0.897	Fbxw5	F-box and WD-40 domain protein 5	
96126_at	20397	-2.3	-2.4	-2.0	-2.2 $\pm$ 0.2	0.704	Sgpl1	Sphingosine phosphate lyase 1	
94405_at	21366	-2.7	-1.5	-1.7	-2.0 $\pm$ 0.2	0.444	Slc6a6	Solute carrier family 6 (neurotransmitter transporter, taurine), member 6	
Response to external stimulus									
101433_at	15064	-2.4	-2.4	-2.4	-2.4 $\pm$ 0.3	0.918	Mr1	Major histocompatibility complex, class I-related	
104228_at	668701	-2.2	-2.2	-2.6	-2.3 $\pm$ 0.4	0.865	EG668701	Similar to Rap guanine nucleotide exchange factor 2	
103250_at	54722	-2.7	-2.1	-2.0	-2.3 $\pm$ 0.2	0.838	Dfna5h	Deafness, autosomal dominant 5 homolog	
104372_at	64296	-2.6	-2.2	-1.9	-2.2 $\pm$ 0.2	0.793	Abhd8	Abhydrolase domain containing 8	
99349_at	16171	-1.7	-2.5	-2.1	-2.1 $\pm$ 0.3	0.681	Il17a	Interleukin 17A	
93218_at	20947	-2.3	-2.0	-1.7	-2.0 $\pm$ 0.4	0.573	Swap70	SWAP complex protein	
Protein modification									
99643_f_at	12876	-3.7	-3.8	-2.7	-3.4 $\pm$ 0.4	0.949	Cpe	Carboxypeptidase E	b
161848_r_at	19260	-2.5	-2.2	-2.4	-2.4 $\pm$ 0.4	0.870	Plpn22	Protein tyrosine phosphatase, non-receptor type 22 (lymphoid)	
92427_at	21812	-2.4	-1.9	-2.3	-2.2 $\pm$ 0.4	0.729	Tgfb1	Transforming growth factor, beta receptor I	
99642_i_at	12876	-3.0	-2.1	-1.3	-2.2 $\pm$ 0.7	0.618	Cpe	Carboxypeptidase E	
100427_at	19277	-2.0	-1.4	-2.7	-2.0 $\pm$ 0.3	0.552	Ptpro	Protein tyrosine phosphatase, receptor type, O	b, c
Transcription from Pol II promoter									
104591_g_at	17260	-2.4	-2.9	-2.5	-2.6 $\pm$ 0.2	0.932	Mef2c	Myocyte enhancer factor 2C	b, c
104590_at	17260	-2.8	-2.0	-2.2	-2.3 $\pm$ 0.2	0.821	Mef2c	Myocyte enhancer factor 2C	a, b
96171_at	54006	-2.1	-1.6	-2.4	-2.0 $\pm$ 0.2	0.580	Deaf1	Deformed epidermal autoregulatory factor 1 (Drosophila)	b

Table 3. (Continued)

H ₁ : M ϕ $\neq$ M ϕ + <i>B. melitensis</i> = M ϕ + <i>B. neotomae</i> = M ϕ + <i>B. ovis</i>									
Probe set	Entrez gene	SLR ^a			Average $\pm$ SEM	EBarrays probability	Gene symbol	Gene product	Additional statistics ^c
		M ^b	N ^b	O ^b					
Cytoskeleton organization and biogenesis									
95705_s_at	11461	-2.2	-3.1	-2.4	-2.6 $\pm$ 0.5	0.887	Actb	Actin, beta, cytoplasmic	
161981_r_at	14246	-2.6	-1.6	-2.4	-2.2 $\pm$ 0.4	0.693	Flg	Filaggrin	
Metabolism									
94872_at	57319	-2.8	-2.5	-2.8	-2.7 $\pm$ 0.4	0.934	Smpd13a	Sphingomyelin phosphodiesterase, acid-like 3A	
161733_at	59010	-2.4	-2.9	-2.1	-2.5 $\pm$ 0.3	0.919	Sqrdl	Sulfide quinone reductase-like	
101972_at	16541	-2.8	-2.4	-2.5	-2.6 $\pm$ 0.3	0.915	Napsa	Napsin A aspartic peptidase	
99667_at	12862	-2.3	-2.2	-2.5	-2.3 $\pm$ 0.5	0.878	Cox6a2	Cytochrome c oxidase, subunit VI a, polypeptide 2	
103538_at	21386	-1.8	-2.1	-2.5	-2.1 $\pm$ 0.3	0.680	Tbx3	T-box 3	
96035_at	12039	-1.7	-2.0	-2.4	-2.0 $\pm$ 0.4	0.606	Bckdha	Branched chain ketoacid dehydrogenase E1, alpha polypeptide	
97560_at	19156	-1.4	-3.0	-1.7	-2.0 $\pm$ 0.4	0.381	Psap	Prosaposin	
Cell communication									
160932_at	17973	-2.6	-2.6	-2.6	-2.6 $\pm$ 0.1	0.964	Nck1	Non-catalytic region of tyrosine kinase adaptor protein 1	a, b
97768_at	13506	-2.2	-2.1	-2.6	-2.3 $\pm$ 0.3	0.803	Dsc2	Desmocollin 2	
Cell growth and/or maintenance									
92695_at	14296	-2.0	-1.6	-2.7	-2.1 $\pm$ 0.3	0.658	Frat1	Frequently rearranged in advanced T-cell lymphomas	
103534_at	15130	-2.8	-2.0	-1.5	-2.1 $\pm$ 0.2	0.626	Hbb-b2	Hemoglobin, beta adult minor chain	b
93736_at	21452	-2.0	-2.0	-2.3	-2.1 $\pm$ 0.3	0.606	Tcn2	Transcobalamin 2	
Miscellaneous classification									
96494_at	75785	-2.9	-2.7	-2.5	-2.7 $\pm$ 0.2	0.959	Klhl24	Kelch-like 24 (<i>Drosophila</i>)	
103933_at	22619	-2.9	-1.9	-2.8	-2.5 $\pm$ 0.2	0.895	Siae	Sialic acid acetyltransferase	b
104299_at	224454	-3.0	-1.9	-2.5	-2.5 $\pm$ 0.3	0.852	Zdhc14	Zinc finger, DHHC domain containing 14	
94299_at	69654	-2.5	-2.0	-2.2	-2.2 $\pm$ 0.2	0.788	Dctn2	Dynactin 2	
162116_r_at	116891	-3.0	-2.7	-1.7	-2.5 $\pm$ 0.9	0.741	Derl2	Der1-like domain family, member 2	
160934_s_at	73094	-3.0	-2.6	-1.6	-2.4 $\pm$ 0.5	0.692	Sgip1	SH3-domain GRB2-like (endophilin) interacting protein 1	
104714_at	105445	-2.2	-1.7	-2.2	-2.0 $\pm$ 0.2	0.611	Dock9	Dedicator of cytokinesis 9	
162075_r_at	17476	-2.7	-1.5	-3.8	-2.7 $\pm$ 0.4	0.551	Mpeg1	Macrophage expressed gene 1	
96464_at	140570	-2.1	-2.2	-1.9	-2.1 $\pm$ 0.2	0.518	Plexnb2	Plexin B2	
160905_s_at	80515	-2.2	-2.2	-2.1	-2.1 $\pm$ 0.3	0.735	A030009	RIKEN cDNA A030009H04 gene	
							H04Rik		
97119_at	99029	-2.6	-2.5	-2.2	-2.4 $\pm$ 0.3	0.911	AI596198	Expressed sequence AI596198	
97752_at	99358	-2.0	-2.1	-2.8	-2.3 $\pm$ 0.3	0.858	E13001	RIKEN cDNA E130013N09 gene	
							3N09Rik		
103748_at	74440	-2.9	-1.6	-2.1	-2.2 $\pm$ 0.2	0.645	4933407	RIKEN cDNA 4933407C03 gene	
							C03Rik		
94069_r_at	67163	-3.4	-2.3	-2.2	-2.6 $\pm$ 0.3	0.946	2610204	RIKEN cDNA 2610204L23 gene	
							L23Rik		

^a Signal log ratio = LOG₂ (experimental *Brucella* spp. infected signal/control uninfected signal) $\pm$ SEM.^b M, N, O indicate *Brucella* spp. used to infect macrophages *B. melitensis*, *B. neotomae* and *B. ovis*, respectively.^c Identified by additional statistical analyses: a, ANOVA $P \leq 0.01$; b, SAM delta = 0.05; c, *B. abortus* microarrays (14).

Table 4A. Differences in Macrophage Transcript Expression when Infected by Different *Brucella* Species

A. <i>B. ovis</i> and <i>B. neotomae</i> Infected Macrophages Express Transcripts Different from <i>B. melitensis</i> or Uninfected Macrophages									
H2: $M\phi \neq M\phi + B. melitensis \neq M\phi + B. neotomae = M\phi + B. ovis$									
Probe set	Entrez gene	SLR ^a				EBarrays probability	Gene symbol	Gene product	Additional statistics ^b
		<i>B. melitensis</i>	<i>B. neotomae</i>	<i>B. ovis</i>					
94337_at	14453	-4.2 ± 0.7	-1.1 ± 0.4	-1.3 ± 0.5	1.000	Gas2	Gas2	Growth arrest specific 2	
162240_r_at	13010	0.8 ± 0.3	3.4 ± 0.2	3.9 ± 0.1	0.999	Cst3	Cst3	Cystatin C	a, b
101793_at	14129	-3.1 ± 0.1	-1.4 ± 0.3	-0.5 ± 0.3	0.903	Fcgr1	Fcgr1	Fc receptor, IgG, high affinity I (FcγRI)	a, b, c
102794_at	12767	-3.6 ± 0.5	-1.4 ± 0.7	-1.7 ± 0.3	0.898	Cxcr4	Cxcr4	Chemokine (C-X-C motif) receptor 4	
101070_at	54484	-3.6 ± 0.5	-1.6 ± 0.6	-1.7 ± 0.3	0.885	Mkm1	Mkm1	Makorin, ring finger protein, 1	
161878_r_at	52250	-2.2 ± 0.8	0.5 ± 0.1	0.3 ± 0.2	0.852	Reep1	Reep1	Receptor accessory protein 1	
93569_f_at	545005	0.6 ± 0.3	2.6 ± 0.4	3.3 ± 0.7	0.829	LOC544988	LOC544988	Similar to Spetex-2F protein	
99377_at	18249	2.7 ± 0.6	0.5 ± 0.9	-0.6 ± 0.6	0.781	Obp1a	Obp1a	Odorant binding protein la	
94240_i_at	19944	1.7 ± 0.9	-0.8 ± 1.3	-0.8 ± 1.0	0.734	Rpl29	Rpl29	Ribosomal protein L29	
104697_at	80837	-3.0 ± 0.7	-1.2 ± 0.9	-0.8 ± 0.4	0.622	Rhoj	Rhoj	ras homolog gene family, member J	
92857_at	19934	-3.3 ± 0.7	-1.2 ± 0.7	-2.0 ± 0.9	0.571	Rpl22	Rpl22	Ribosomal protein L22	
102706_i_at	16625	-2.7 ± 0.9	-0.6 ± 0.4	-0.1 ± 0.3	0.566	Serpina3c	Serpina3c	Serine (or cysteine) proteinase inhibitor, clade A, member 3C	

Table 4B.

B. <i>B. melitensis</i> and <i>B. neotomae</i> Infected Macrophages Express Transcripts Different from <i>B. ovis</i> or Uninfected Macrophages									
H3: $M\phi \neq M\phi + B. melitensis = M\phi + B. neotomae \neq M\phi + B. ovis$									
Probe set	Entrez gene	SLR ^a				EBarrays probability	Gene symbol	Gene product	
		<i>B. melitensis</i>	<i>B. neotomae</i>	<i>B. ovis</i>					
161420_r_at	13478	-0.3 ± 0.7	-0.2 ± 1.3	-3.2 ± 0.5	0.993	Dpagt1	Dpagt1	Dolichyl-phosphate acetylgluco-saminephosphotransferase 1 (GlcNAc-1-P transferase)	
160797_r_at	12412	-2.1 ± 0.4	-2.0 ± 0.4	0.6 ± 0.2	0.975	Cbx1	Cbx1	Chromobox homolog 1 (Drosophila HP1 beta)	
92600_f_at	13117	-2.4 ± 0.6	-2.4 ± 0.7	0.0 ± 0.1	0.953	Cyp4a10	Cyp4a10	Cytochrome P450, family 4, subfamily a, polypeptide 10	
161128_r_at	544834	0.5 ± 1.3	1.8 ± 0.9	3.7 ± 0.8	0.9361	6030426L16Rik	6030426L16Rik	Similar to Hippocalcin-like protein 1, Visinin-like protein 3	
99885_at	16019	0.6 ± 0.9	1.0 ± 0.3	3.1 ± 0.6	0.738	Igh-6	Igh-6	Immunoglobulin heavy chain 6 (heavy chain of IgM)	
93566_at	74246	-1.5 ± 0.2	-2.6 ± 0.6	0.1 ± 0.3	0.711	Gale	Gale	Galactose-4-epimerase, UDP	
162076_r_at	14800	-2.2 ± 0.7	-3.0 ± 0.5	-0.5 ± 0.4	0.681	Gria2	Gria2	Glutamate receptor, ionotropic, AMPA2 (alpha 2)	
160728_r_at	73945	-2.1 ± 1.1	-0.4 ± 0.6	-4.4 ± 0.6	0.6579	Ottd4	Ottd4	OTU domain containing 4	
AFFX-BioDn-3_st	1037747	-1.6 ± 0.3	-0.4 ± 0.1	1.4 ± 0.1	0.6294	—	—	Biotin synthesis protein bioC	
161131_r_at	228889	-0.6 ± 0.4	-0.7 ± 0.1	-2.7 ± 0.1	0.5167	—	—	Dead (asp-glu-ala-asp) box polypeptide 27	
102299_at	18750	-1.8 ± 0.7	-1.2 ± 0.5	-3.3 ± 0.4	0.508	Prkca	Prkca	Protein kinase C, alpha	

Table 4C.

C. <i>B. melitensis</i> and <i>B. ovis</i> Infected Macrophages Express Transcripts Different from <i>B. neotomae</i> or Uninfected Macrophages									
H ₄ : Mφ ≠ Mφ + <i>B. melitensis</i> = Mφ + <i>B. ovis</i> ≠ Mφ + <i>B. neotomae</i>									
Probe set	Entrez gene	SLR ^a			EBarrays probability	Gene symbol	Gene product	Additional statistics ^b	
		<i>B. melitensis</i>	<i>B. neotomae</i>	<i>B. ovis</i>					
94051_at	13612	3.0 ± 0.4	0.5 ± 0.4	3.2 ± 0.3	0.998	Edil3	EGF-like repeats and discordin 1-like domains 3	a, b	
96785_at	80880	-3.2 ± 0.6	-0.5 ± 1.1	-2.7 ± 0.7	0.994	Ankrd47	Ankyrin repeat domain 47		
92913_at	26874	-4.1 ± 0.3	-2.0 ± 0.4	-3.9 ± 0.3	0.920	Abcd2	ATP-binding cassette, sub-family D (ALD), member 2	a, b	
103467_g_at	54151	-2.3 ± 0.3	-0.6 ± 0.3	-2.9 ± 0.3	0.897	Cyhr1	Cysteine and histidine rich 1		
94755_at	16175	3.9 ± 0.4	1.7 ± 0.7	3.1 ± 0.4	0.693	Il1a	Interleukin 1 alpha	b, c	
93568_i_at		1.4 ± 0.0	-1.1 ± 0.7	1.2 ± 0.5	0.556	LOC544988	Similar to Spetex-2F protein		
92504_at	15574	2.1 ± 0.6	-0.1 ± 0.4	1.8 ± 0.7	0.501	Hus1	Hus1 homolog (S. pombe)		
104592_i_at	17260	-3.7 ± 0.2	-1.9 ± 0.3	-3.3 ± 0.2	0.473	Mef2c	Myocyte enhancer factor 2C	a, b	
102798_at	11535	0.9 ± 0.6	2.4 ± 0.6	-0.3 ± 0.7	0.444	Adm	Adrenomedullin		
102374_at	53902	-2.9 ± 0.3	-1.0 ± 0.4	-2.2 ± 0.5	0.425	Dscr12	Down syndrome critical region gene 1-like 2	a, b	

^a Signal log ratio = LOG₂ (experimental *Brucella* spp. infected signal/control uninfected signal) ± SEM.

^b Identified by additional statistical analyses: a, ANOVA $P \leq 0.01$; b, = SAM delta = 0.05; c, *B. abortus* microarrays (14).

murine macrophages in contrast to *in vivo* human infection where *B. neotomae* is unable to initiate or maintain infection (29, 30). The IRF-1^{-/-} mouse model can distinguish degrees of *Brucella* virulence (5), where *B. melitensis* and *B. neotomae* induce death by 9–12 days (6), while *B. ovis* does not (E. Petersen, Personal Communication). The differing responses between rough (*B. ovis*) and smooth (*B. melitensis* and *B. neotomae*) *Brucella* in mice led us to explore transcription profiles of murine macrophages following infection by these three *Brucella* strains. A comparison of host response among infections by differing *Brucella* spp. may identify common responses to infection, as well as distinguish genes and pathways specific to each *Brucella* spp. infection. Identifying alterations in the macrophage transcriptome may provide greater understanding of host mechanisms involved in pathogen killing and bacterial regulation.

Common Macrophage Transcriptional Response by *B. melitensis*, *B. neotomae* and *B. ovis* Infections. At 4 hr, low numbers of intracellular smooth *B. melitensis* and *B. neotomae* were observed, as reported by others, prior to *Brucella* reaching its replicative niche (22, 31, 32). However, RAW 264.7 macrophage uptake of rough *B. ovis* after 4 hr of infection was almost two logarithmic units higher than smooth *B. melitensis* or *B. neotomae*, perhaps from non-lipid raft mediated phagocytosis, although this is untested for rough *B. ovis* (22, 31, 32). Yet, microarray analysis of rough *B. ovis*-infected macrophages was qualitatively similar to those from smooth *B. melitensis* and *B. neotomae* infections. This similarity suggests that macrophages sense and respond to different *Brucella* species by a common profile of transcriptional changes.

The majority of these common response genes were inflammatory related suggesting similar activating and/or regulating effects of *Brucella* on the macrophage regardless of the *Brucella* species. Host response to *Brucella* spp. encompassed pro-inflammation independent of infection differences between rough and smooth *Brucella* spp. In comparing the common macrophage transcription profile to differing *Brucella* spp., we hypothesized that a large number of host genes with overlapping gene ontology function would be identified similar to that observed in wild-type and mutant *Salmonella* (33). Others have reported that LPS and CpG DNA can activate TLR signaling activating NFκB and AP-1 pathways inducing innate immune mechanisms (34). Adaptive immune mechanisms may be activated by Schlafen 2 (*Slfn2*) expression that regulates T cell activity as well as innate immunity (34). Pro-inflammatory gene expression may reflect a host protective response as many genes were chemotaxis related, potentially recruiting additional macrophages and immune cells to the site of *Brucella* spp. infected macrophages.

Genes that could serve to subdue or counteract inflammatory response genes, however, were also observed




Figure 2. Transcript level changes in select genes following *Brucella* spp. infection of RAW 264.7 (Panel A) or J774A.1 (Panel B) macrophages. RT-PCR results confirm genes with increased A or C or decreased B or D transcript levels following infection of RAW 264.7 or J774A.1 cells, respectively. Each RNA sample was normalized using *Hprt* transcript levels. Signal log ratios (SLR), a comparison of each infected sample to uninfected control RAW 264.7 samples, were averaged from four independent RNA isolations and error bars represent the SEM.

as upregulated in macrophages. For example, increased expression of *Socs3* was observed. *Socs3* participates in a negative feedback loop of cytokine signaling limiting the extent of cytokine in innate and adaptive immune responses (35). Also, *IL-13R* and *Gadd45b* were increased in transcription suggesting regulation of activated macrophages. Transcription of *Bcl2*, *Bat4* and *Phlda1* that participate in blocking apoptotic mechanisms was upregulated. In agreement with this finding, smooth *Brucella* spp. can protect against apoptosis (36), while engineered rough *Brucella* spp. can induce apoptosis (37). Consequently, the upregulation of genes that block apoptotic mechanisms may oppose upregulated inflammatory genes to camouflage immune recognition and produce a protective environment for *Brucella* replication. Also, decreased transcript levels of *Psap* (lysosomal catabolic enzyme), *Sgip3* (clathrin mediated endocytosis) and *Mpeg1* (a perforin like protein) would suggest that *Brucella* induces evasion strategies to protect its intracellular niche. Others have suggested evasion strategies such as *Brucella* altering host cell responses by shielding PAMPs from Toll-like receptors (38) and LPS masking of MHC molecules (39).

Of the macrophage genes commonly upregulated by all *Brucella* species tested, first, eicosanoids are important in mammalian repair (40). In macrophages infected with *Brucella* we observed induction of eicosanoid *Ptgs2*, also

termed *Cox2*, encoding the cyclooxygenase 2 enzyme. Eicosanoids are considered important in antigen presentation and in macrophage effector functions, and play an important role in contributing to inflammation (40).

A second common inflammatory signal elicited by *Brucella* infection of macrophages is *Gadd45*. *Gadd45A* is an 18 kDa acidic nuclear protein involved in maintenance of genomic stability, DNA repair, cell cycle checkpoints and suppression of cell growth. *Gadd45* is also a known stress response gene with the ability to regulate MAP-kinase signaling (41). Regulation of MAP-kinase signaling by *Gadd45* can affect response to extracellular stimuli and regulate gene expression, mitosis, differentiation and cell survival/apoptosis. *GADD45A* may therefore be expressed as a stress response protein to inhibit macrophage growth during *Brucella* infection.

A third common inflammatory signal elicited by *Brucella* infection is an odorant binding protein. Odorant binding proteins (OBPs) belong to the lipocalin family of proteins involved in extracellular transport of hydrophobic ligands. OBPs were originally identified in nasal mucus and mucosa of insects and were proposed as molecular shuttles between the air mucus interface and the olfactory receptor binding sites. Their function in mammals is unknown, but OBPs can bind odorants of diverse chemical structures with a higher affinity for aldehydes and large fatty acids (42). In




Figure 3. Induction of TNF α by the *B. melitensis* in BMDMs from C57BL/6 mice. Cytokine levels were assayed from supernatants of *B. melitensis* 16M infected or medium alone bone-marrow derived macrophages (BMDMs). BMDMs were cultured 5–7 days pre-infection. Cytokines were measured in pg/mL by ELISA at 12 hr post-infection. Data are from 7 mice/experiment.

macrophages, OBPs may serve as chaperones for signaling molecules.

Macrophage Transcriptome Response to *B. melitensis* Differ from *B. neotomae* and *B. ovis*.

Macrophages infected with *B. melitensis* elicited a specific response with altered transcription in twelve genes compared to *B. neotomae* and *B. ovis* (H₂) (Table 4, part A). These genes provide insight into macrophage mechanisms used by the various *Brucella* species to establish a replicative niche. For example, two genes with established function during inflammation, *FcγRI* (43) and *Cxcr4* (44), have decreased expression in *B. melitensis* infected macrophages compared to *B. neotomae* and *B. ovis* infections. *FcγRI* (CD64) may be involved in regulating antigen presentation by limiting additional phagocytosis and committing the cell to present antigens of engulfed pathogens (45). Decreased levels of *FcγRI* may be advantageous for bacteria by reducing downstream signaling initiation, including antibody dependent cell mediated cytotoxicity (ADCC) (46), production of reactive oxygen and nitrogen species (47, 48), and increased phagocytosis and cytokine secretion (49). Downregulation of *Cxcr4* transcription in combination with *Socs3* upregulation may reduce macrophage chemotaxis, adhesion and other inflammatory responses (50, 51). Suppressing several immune pathways initiated at the cell membrane, virulent *Brucella* may secure an intracellular replicative niche.

Three other genes with decreased expression in macrophages infected with *B. melitensis* compared to *B. neotomae* and *B. ovis* (H₂ hypothesis) are endopeptidase inhibitors, *cystatin C* and *serpina3c* and the actin associated protein encoded by *Gas2*. These natural peptidase inhibitors may influence cell migration, chemotaxis, proliferation, phagocytosis and respiratory burst (52–55). Interestingly, the treatment of mice harboring *Leishmania* with cystatin and a sub-optimal dose of IFN- γ led to parasite clearance and shifted the non-productive Th2 (type 2, humoral)

response towards a Th1 response (56). Nitric oxide (NO) production (57) and phagosomal processing (55) are mechanisms where these endopeptidase inhibitors act and modulate intracellular infection. These same mechanisms are vital to progression of *Brucella* spp. infection (58–60) and exogenous modulation may provide the immunologic boost necessary for pathogen clearance. *Brucella* spp. may specifically alter the protease-inhibitor balance by bacterial production of extracellular proteases and cleavage of host endopeptidase inhibitors, similar to how *Staphylococcus aureus* functions in accessing nutrients and facilitating dissemination (61). Lastly, *Gas2* was specifically down-regulated in *B. melitensis*. GAS2-like protein 1 is an actin-associated protein expressed at high levels in growth-arrested macrophages (62). Inhibition of *Gas2* transcription may increase macrophage survival and growth, an advantage for *Brucella* spp. survival within an intracellular niche.

Eleven genes were altered in expression (Table 4, part B) by *B. melitensis* and *B. neotomae* but not by *B. ovis* (H₃). These genes could not be grouped according to function, but did categorize the *Brucella* spp. according to LPS type, rough or smooth. While *Brucella* spp. uptake based on smooth or rough phenotype did not affect transcript levels of many inflammatory-related genes in our study (H₁), LPS may subtly affect transcription when comparing *B. melitensis* and *B. neotomae* to *B. ovis*. Microarray analyses of macrophages responding to various bacteria indicate that an early response to infection parallels the response to LPS alone and is conserved across bacterial genera (63–65). For example, macrophage response to *Salmonella* versus purified *Salmonella* LPS is similar (64). However, *Brucella* LPS induces a 100-fold less stimulation of macrophages than enterobacterial LPS (2, 66), and may have less effect on macrophage transcription.

Cytochrome P450 4a10 (*Cyp4a10*) and *protein kinase C- α* (*Prkca*) may provide insight into intracellular pathways that differ between *Brucella* species (H₃) (Table 4, part B). A wide variety of bacterial and parasitic infections depress expression of *cytochrome P450* (67), a gene important in detoxification (68). *Cyp4a10* can enhance oxidative stress, through production of reactive oxygen species and reactive nitrogen species (69); therefore, decreased transcription induced by bacteria with smooth LPS could protect intracellular *Brucella* from innate killing mechanisms. Another gene with oxidative related functions, *Prkca*, has decreased transcript levels in the infected macrophage. *Prkca* is important in regulating phagosome-lysosome fusion (70), intracellular vesicle regulation (71) and generation of reactive oxygen species (72), events that may be regulated differently by diverse *Brucella* spp. Constitutive knock down of *Prkca* during intracellular infection with *Legionella*, *Leishmania* and *Salmonella* (48, 73) leads to increased pathogen survival, likely through suppression of host respiratory burst. Downregulation of *Cyp4a10* and *Prkca* pathways may prevent additional oxidative stress in the macrophage during smooth *Brucella*

spp. infection, while *Brucella* spp. production of Cu, Zn superoxide dismutase (SodC) protects *Brucella* from oxidative damage (74). These results suggest that down-regulation of *Cyp4a10* and *Pkrca* may shift the macrophage toward a less hostile intracellular environment for *Brucella*.

Finally altered transcription was identified when *B. melitensis* and *B. ovis* were considered dissimilar from *B. neotomae* (H₄). Nine of ten genes had decreased transcript levels in *B. neotomae* infection compared to the other *Brucella* spp., the most important examples being *IL-1a*, *Abcd2* and *mef2c*. *B. neotomae* may moderate the IL-1 α transcription earlier or through alternative stimulation. *Abcd2* knock down has been correlated with oxidative stress or tissue damage (75) and abnormalities in mitochondria, Golgi and endoplasmic reticulum of host cells (76). The importance of *Abcd* transporters is reinforced by decreased transcript levels of other *Abcd* family members during *Brucella* infection (9, 77). Lastly, the alteration of *mef2c* expression suggests that *B. neotomae* infection may render RAW 264.7 macrophages more susceptible to activation-induced cell death (78) than *B. melitensis* or *B. ovis* infections. When compared to *B. melitensis* and *B. ovis*, *B. neotomae* infected macrophages appeared to have decreased immuno-stimulatory capability resulting in reduced oxidative stress and increased susceptibility to cell death.

Brucella infection of RAW 264.7 and J774A.1 macrophage cell lines resulted in similar results. While RAW 264.7 and J774A.1 cells are macrophage-like in lysozyme secretion (79), phagocytosis of particles, response to LPS stimulus (80) and activation state (81), differences do exist in the abilities of these cells to mediate cytolytic activity (81), synthesize nitrite and nitrate (80), and potentially other macrophage trafficking and defense mechanisms. Our microarray results were supported by similar transcription profiles between these *Brucella* infected macrophage cell lines. However, internalization of smooth *Brucella* by phagocytic macrophages occurs faster than by nonphagocytic cells such as trophoblasts (82) and may contribute to differences in microarray results between phagocytic and nonphagocytic cells. Ultimately, individual genes of interest must be further investigated for influence on *Brucella* spp. infections. In conclusion, the present study evaluated the macrophage transcriptome altered during *B. melitensis*, *B. neotomae* and *B. ovis* infections. Common changes in gene expression were observed among *Brucella* spp. infected macrophages suggesting similar innate immune mechanisms by macrophages when responding to different *Brucella* species. These changes were related to increased chemokine, alteration in cytokine signals and defense responses. Also, these three *Brucella* spp. induced decreased transcription of select genes, suggesting that repressed transcription may result from pathogen-specific manipulations. Further, a subset of macrophage genes was differentially altered based on a particular *Brucella* species that may reflect the differing intracellular survival abilities of each

Brucella species. Microarray analysis of host transcription provides a foundation to understand variations in *Brucella* spp. infection of mice. Future studies will reveal variations in the intracellular survival strategies used by different *Brucella* spp. by evaluating the contribution of individual host genes during infection.

The authors would like to thank S. Splinter and W. Nelson at the Genome Center of Wisconsin for GeneChip® hybridization and scanning, C. Kendzierski for assistance with EBarrays statistical analysis and E. Petersen for personal communications.

1. Woods JB, Ed. USAMRIID's Medical Management of Biological Casualties Handbook (6th ed.). Fort Detrick, MD: U.S. Army Medical Research Institute of Infectious Diseases, 2005.
2. Baldwin CL, Goenka R. Host immune responses to the intracellular bacteria *Brucella*: does the bacteria instruct the host to facilitate chronic infection? *Crit Rev Immunol* 26:407–442, 2006.
3. Halling SM, Peterson-Burch BD, Bricker BJ, Zuermer RL, Qing Z, Li LL, Kapur V, Alt DP, Olsen SC. Completion of the genome sequence of *Brucella abortus* and comparison to the highly similar genomes of *Brucella melitensis* and *Brucella suis*. *J Bacteriol* 187:2715–2726, 2005.
4. Rajashekara G, Glasner JD, Glover DA, Splitter GA. Comparative whole-genome hybridization reveals genomic islands in *Brucella* species. *J Bacteriol* 186:5040–5051, 2004.
5. Ko J, Gendron-Fitzpatrick A, Ficht TA, Splitter GA. Virulence criteria for *Brucella abortus* strains as determined by interferon regulatory factor 1-deficient mice. *Infect Immun* 70:7004–7012, 2002.
6. Baek SH, Rajashekara G, Splitter GA, Shapleigh JP. Denitrification genes regulate *Brucella* virulence in mice. *J Bacteriol* 186:6025–6031, 2004.
7. Eskra L, Mathison A, Splitter G. Microarray analysis of mRNA levels from RAW264.7 macrophages infected with *Brucella abortus*. *Infect Immun* 71:1125–1133, 2003.
8. Celli J, de Chastellier C, Franchini DM, Pizarro-Cerda J, Moreno E, Gorvel JP. *Brucella* evades macrophage killing via VirB-dependent sustained interactions with the endoplasmic reticulum. *J Exp Med* 198: 545–556, 2003.
9. He Y, Reichow S, Ramamoorthy S, Ding X, Lathigra R, Craig JC, Sobral BW, Schurig GG, Sriranganathan N, Boyle SM. *Brucella melitensis* triggers time-dependent modulation of apoptosis and down-regulation of mitochondrion-associated gene expression in mouse macrophages. *Infect Immun* 74:5035–5046, 2006.
10. Caron E, Liautard JP, Kohler S. Differentiated U937 cells exhibit increased bactericidal activity upon LPS activation and discriminate between virulent and avirulent *Listeria* and *Brucella* species. *J Leukoc Biol* 56:174–181, 1994.
11. Jiang X, Baldwin CL. Effects of cytokines on intracellular growth of *Brucella abortus*. *Infect Immun* 61:124–134, 1993.
12. Rittig MG, Alvarez-Martinez MT, Porte F, Liautard JP, Rouot B. Intracellular survival of *Brucella* spp. in human monocytes involves conventional uptake but special phagosomes. *Infect Immun* 69:3995–4006, 2001.
13. Eskra L, Canavessi A, Carey M, Splitter G. *Brucella abortus* genes identified following constitutive growth and macrophage infection. *Infect Immun* 69:7736–7742, 2001.
14. Newton MA, Kendzierski CM. Parametric empirical Bayes methods for microarrays. In: Parmigiani G, Garrett ES, Irizarry RA, Zeger SL, Eds. *The Analysis of Gene Expression Data: Methods and Software*. New York: Springer, pp259–276, 2003.
15. Kendzierski CM, Newton MA, Lan H, Gould MN. On parametric

- empirical Bayes methods for comparing multiple groups using replicated gene expression profiles. *Stat Med* 22:3899–3914, 2003.
16. Zar JH. *Biostatistical Analysis*. New Jersey: Prentice Hall, 1999.
 17. Chu G, Narasimhan B, Tibshirani R, Tusher V. SAM “Significance Analysis of Microarrays” Users Guide and Technical Document. Available at: <http://www-stat.stanford.edu/~tib/SAM/index.html>. 2002.
 18. Tusher VG, Tibshirani R, Chu G. Significance analysis of microarrays applied to the ionizing radiation response. *Proc Natl Acad Sci U S A* 98:5116–5121, 2001.
 19. Saeed AI, Bhagabati NK, Braisted JC, Liang W, Sharov V, Howe EA, Li J, Thiagarajan M, White JA, Quackenbush J. TM4 microarray software suite. *Methods Enzymol* 411:134–193, 2006.
 20. Saeed AI, Sharov V, White J, Li J, Liang W, Bhagabati N, Braisted J, Klapa M, Currier T, Thiagarajan M, Sturn A, Snuffin M, Rezantsev A, Popov D, Ryltsov A, Kostukovich E, Borisovsky I, Liu Z, Vinsavich A, Trush V, Quackenbush J. TM4: a free, open-source system for microarray data management and analysis. *Biotechniques* 34:374–378, 2003.
 21. Rozen S, Skaletsky H. Primer3 on the WWW for general users and for biologist programmers. *Methods Mol Biol* 132:365–386, 2000.
 22. Jimenez de Bagues MP, Terraza A, Gross A, Dornand J. Different responses of macrophages to smooth and rough *Brucella* spp.: relationship to virulence. *Infect Immun* 72:2429–2433, 2004.
 23. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* 3:1–12, 2002.
 24. Jenner RG, Young RA. Insights into host responses against pathogens from transcriptional profiling. *Nat Rev Microbiol* 3:281–294, 2005.
 25. Chaves-Olarte E, Guzman-Verri C, Meresse S, Desjardins M, Pizarro-Cerda J, Badilla J, Gorvel JP, Moreno E. Activation of Rho and Rab GTPases dissociates *Brucella abortus* internalization from intracellular trafficking. *Cell Microbiol* 4:663–676, 2002.
 26. Naroeni A, Porte F. Role of cholesterol and the ganglioside GM(1) in entry and short-term survival of *Brucella suis* in murine macrophages. *Infect Immun* 70:1640–1644, 2002.
 27. Radonic A, Thulke S, Mackay IM, Landt O, Siegert W, Nitsche A. Guideline to reference gene selection for quantitative real-time PCR. *Biochem Biophys Res Commun* 313:856–862, 2004.
 28. Dheda K, Huggett JF, Bustin SA, Johnson MA, Rook G, Zumla A. Validation of housekeeping genes for normalizing RNA expression in real-time PCR. *Biotechniques* 37:112–114, 116, 118–119, 2004.
 29. Stoenner HG. The behavior of *Brucella neotomae* and *Brucella suis* in reciprocal superinfection experiments in mice and guinea pigs. *Am J Vet Res* 24:376–380, 1963.
 30. Stoenner HG, Lackman DB. A new species of *Brucella* isolated from the desert wood rat, *Neotoma lepida* Thomas. *Am J Vet Res* 18:947–951, 1957.
 31. Rittig MG, Kaufmann A, Robins A, Shaw B, Sprenger H, Gerns D, Foulongne V, Rouot B, Dornand J. Smooth and rough lipopolysaccharide phenotypes of *Brucella* induce different intracellular trafficking and cytokine/chemokine release in human monocytes. *J Leukoc Biol* 74:1045–1055, 2003.
 32. Porte F, Naroeni A, Ouahrani-Bettache S, Liautard JP. Role of the *Brucella suis* lipopolysaccharide O antigen in phagosomal genesis and in inhibition of phagosome-lysosome fusion in murine macrophages. *Infect Immun* 71:1481–1490, 2003.
 33. Detweiler CS, Cunanan DB, Falkow S. Host microarray analysis reveals a role for the *Salmonella* response regulator phoP in human macrophage cell death. *Proc Natl Acad Sci U S A* 98:5850–5855, 2001.
 34. Sohn WJ, Kim D, Lee KW, Kim MS, Kwon S, Lee Y, Kim DS, Kwon HJ. Novel transcriptional regulation of the *schlafen-2* gene in macrophages in response to TLR-triggered stimulation. *Mol Immunol* 44:3273–3282, 2007.
 35. Dimitriou ID, Clemenza L, Scotter AJ, Chen G, Guerra FM, Rottapel R. Putting out the fire: coordinated suppression of the innate and adaptive immune systems by SOCS1 and SOCS3 proteins. *Immunol Rev* 224:265–283, 2008.
 36. Gross A, Terraza A, Ouahrani-Bettache S, Liautard JP, Dornand J. In vitro *Brucella suis* infection prevents the programmed cell death of human monocytic cells. *Infect Immun* 68:342–351, 2000.
 37. Fernandez-Prada CM, Zelazowska EB, Nikolich M, Hadfield TL, Roop RM 2nd, Robertson GL, Hoover DL. Interactions between *Brucella melitensis* and human phagocytes: bacterial surface O-Polysaccharide inhibits phagocytosis, bacterial killing, and subsequent host cell apoptosis. *Infect Immun* 71:2110–2119, 2003.
 38. Rajashekara G, Covert J, Petersen E, Eskra L, Splitter G. Genomic island 2 of *Brucella melitensis* is a major virulence determinant: functional analyses of genomic islands. *J Bacteriol* 190:6243–6252, 2008.
 39. Lapaque N, Forquet F, de Chastellier C, Mishal Z, Jolly G, Moreno E, Moriyon I, Heuser JE, He HT, Gorvel JP. Characterization of *Brucella abortus* lipopolysaccharide macrodomains as mega rafts. *Cell Microbiol* 8:197–206, 2006.
 40. Harizi H, Gualde N. The impact of eicosanoids on the crosstalk between innate and adaptive immunity: the key roles of dendritic cells. *Tissue Antigens* 65:507–514, 2005.
 41. Zhan Q. Gadd45a, a p53- and BRCA1-regulated stress protein, in cellular response to DNA damage. *Mutat Res* 569:133–143, 2005.
 42. Briand L, Eloit C, Nespoulous C, Bezirard V, Huet JC, Henry C, Blon F, Trotier D, Pernollet JC. Evidence of an odorant-binding protein in the human olfactory mucus: location, structural characterization, and odorant-binding properties. *Biochemistry* 41:7241–7252, 2002.
 43. Nakamura K, Malykhin A, Coggeshall KM. The Src homology 2 domain-containing inositol 5-phosphatase negatively regulates Fcγ receptor-mediated phagocytosis through immunoreceptor tyrosine-based activation motif-bearing phagocytic receptors. *Blood* 100:3374–3382, 2002.
 44. Wragg A, Mellad JA, Beltran LE, Konoplyannikov M, San H, Boozer S, Deans RJ, Mathur A, Lederman RJ, Kovacic JC, Boehm M. VEGFR1/CXCR4-positive progenitor cells modulate local inflammation and augment tissue perfusion by a SDF-1-dependent mechanism. *J Mol Med* 86:1221–1232, 2008.
 45. Boldrick JC, Alizadeh AA, Diehn M, Dudoit S, Liu CL, Belcher CE, Botstein D, Staudt LM, Brown PO, Relman DA. Stereotyped and specific gene expression programs in human innate immune responses to bacteria. *Proc Natl Acad Sci U S A* 99:972–977, 2002.
 46. Zheng L, Zomerdijs TP, Aarnoudse C, van Furth R, Nibbering PH. Role of protein kinase C isozymes in Fcγ receptor-mediated intracellular killing of *Staphylococcus aureus* by human monocytes. *J Immunol* 155:776–784, 1995.
 47. Joshi T, Butchar JP, Tridandapani S. Fcγ receptor signaling in phagocytes. *Int J Hematol* 84:210–216, 2006.
 48. Dasgupta D, Chakraborty P, Basu MK. Ligation of Fc receptor of macrophages stimulates protein kinase C and anti-leishmanial activity. *Mol Cell Biochem* 209:1–8, 2000.
 49. Loegering DJ, Lennartz MR. Signaling pathways for Fcγ receptor-stimulated tumor necrosis factor-α secretion and respiratory burst in RAW 264.7 macrophages. *Inflammation* 28:23–31, 2004.
 50. Paranaivitana CM, Zelazowska E, Das R, Izadjoo M, Jett M, Hoover D. Identification of novel genes in the memory response to *Brucella* infection by cDNA arrays. *Mol Cell Probes* 19:341–348, 2005.
 51. Soriano SF, Hernanz-Falcon P, Rodriguez-Frade JM, De Ana AM, Garzon R, Carvalho-Pinto C, Vila-Coro AJ, Zaballos A, Balomenos D, Martinez AC, Mellado M. Functional inactivation of CXCR4 chemokine receptor 4-mediated responses through SOCS3 up-regulation. *J Exp Med* 196:311–321, 2002.
 52. Hiemstra PS. Novel roles of protease inhibitors in infection and inflammation. *Biochem Soc Trans* 30:116–120, 2002.

53. Vray B, Hartmann S, Hoebeke J. Immunomodulatory properties of cystatins. *Cell Mol Life Sci* 59:1503–1512, 2002.
54. Forsyth S, Horvath A, Coughlin P. A review and comparison of the murine alpha1-antitrypsin and alpha1-antichymotrypsin multigene clusters with the human clade A serpins. *Genomics* 81:336–345, 2003.
55. Lefebvre C, Cocquerelle C, Vandenbulcke F, Hot D, Huot L, Lemoine Y, Salzet M. Transcriptomic analysis in the leech *Theromyzon tessulatatum*: involvement of cystatin B in innate immunity. *Biochem J* 380:617–625, 2004.
56. Das L, Datta N, Bandyopadhyay S, Das PK. Successful therapy of lethal murine visceral leishmaniasis with cystatin involves up-regulation of nitric oxide and a favorable T cell response. *J Immunol* 166:4020–4028, 2001.
57. Mukherjee S, Ukil A, Das PK. Immunomodulatory peptide from cystatin, a natural cysteine protease inhibitor, against leishmaniasis as a model macrophage disease. *Antimicrob Agents Chemother* 51:1700–1707, 2007.
58. Ko J, Gendron-Fitzpatrick A, Splitter GA. Susceptibility of IFN regulatory factor-1 and IFN consensus sequence binding protein-deficient mice to brucellosis. *J Immunol* 168:2433–2440, 2002.
59. Wang M, Qureshi N, Soeurt N, Splitter G. High levels of nitric oxide production decrease early but increase late survival of *Brucella abortus* in macrophages. *Microb Pathog* 31:221–230, 2001.
60. Porte F, Liautard JP, Kohler S. Early acidification of phagosomes containing *Brucella suis* is essential for intracellular survival in murine macrophages. *Infect Immun* 67:4041–4047, 1999.
61. Vincents B, Onnerfjord P, Gruca M, Potempa J, Abrahamson M. Down-regulation of human extracellular cysteine protease inhibitors by the secreted staphylococcal cysteine proteases, staphopain A and B. *Biol Chem* 388:437–446, 2007.
62. Stramer B, Winfield M, Shaw T, Millard TH, Woolner S, Martin P. Gene induction following wounding of wild-type versus macrophage-deficient *Drosophila* embryos. *EMBO Rep* 9:465–471, 2008.
63. Nau GJ, Richmond JF, Schlesinger A, Jennings EG, Lander ES, Young RA. Human macrophage activation programs induced by bacterial pathogens. *Proc Natl Acad Sci U S A* 99:1503–1508, 2002.
64. Rosenberger CM, Scott MG, Gold MR, Hancock RE, Finlay BB. *Salmonella typhimurium* infection and lipopolysaccharide stimulation induce similar changes in macrophage gene expression. *J Immunol* 164:5894–5904, 2000.
65. Huang H, Park CK, Ryu JY, Chang EJ, Lee Y, Kang SS, Kim HH. Expression profiling of lipopolysaccharide target genes in RAW264.7 cells by oligonucleotide microarray analyses. *Arch Pharm Res* 29:890–897, 2006.
66. Duenas AI, Orduna A, Crespo MS, Garcia-Rodriguez C. Interaction of endotoxins with Toll-like receptor 4 correlates with their endotoxic potential and may explain the proinflammatory effect of *Brucella* spp. LPS. *Int Immunol* 16:1467–1475, 2004.
67. Renton KW. Cytochrome P450 regulation and drug biotransformation during inflammation and infection. *Curr Drug Metab* 5:235–243, 2004.
68. Hossain H, Tchatalbachev S, Chakraborty T. Host gene expression profiling in pathogen-host interactions. *Curr Opin Immunol* 18:422–429, 2006.
69. Morgan ET. Regulation of cytochrome p450 by inflammatory mediators: why and how? *Drug Metab Dispos* 29:207–212, 2001.
70. Ng Yan Hing JD, Desjardins M, Descoteaux A. Proteomic analysis reveals a role for protein kinase C-alpha in phagosome maturation. *Biochem Biophys Res Commun* 319:810–816, 2004.
71. Chen YW, Lang ML, Wade WF. Protein kinase C-alpha and -delta are required for FcalphaR (CD89) trafficking to MHC class II compartments and FcalphaR-mediated antigen presentation. *Traffic* 5:577–594, 2004.
72. von Knethen A, Tautenhahn A, Link H, Lindemann D, Brune B. Activation-induced depletion of protein kinase C alpha provokes desensitization of monocytes/macrophages in sepsis. *J Immunol* 174:4960–4965, 2005.
73. St-Denis A, Caouras V, Gervais F, Descoteaux A. Role of protein kinase C-alpha in the control of infection by intracellular pathogens in macrophages. *J Immunol* 163:5505–5511, 1999.
74. Gee JM, Valderas MW, Kovach ME, Grippe VK, Robertson GT, Ng WL, Richardson JM, Winkler ME, Roop RM 2nd. The *Brucella abortus* Cu,Zn superoxide dismutase is required for optimal resistance to oxidative killing by murine macrophages and wild-type virulence in experimentally infected mice. *Infect Immun* 73:2873–2880, 2005.
75. Lu JF, Barron-Casella E, Deering R, Heinzer AK, Moser AB, deMesy Bentley KL, Wand GS, McGuinness MC, Pei Z, Watkins PA, Pujol A, Smith KD, Powers JM. The role of peroxisomal ABC transporters in the mouse adrenal gland: the loss of Abcd2 (ALDR), Not Abcd1 (ALD), causes oxidative damage. *Lab Invest* 87:261–272, 2007.
76. Wanders RJ, Visser WF, van Roermund CW, Kemp S, Waterham HR. The peroxisomal ABC transporter family. *Pflugers Arch* 453:719–734, 2007.
77. Roux CM, Rolan HG, Santos RL, Beremand PD, Thomas TL, Adams LG, Tsois RM. *Brucella* requires a functional Type IV secretion system to elicit innate immune responses in mice. *Cell Microbiol* 9:1851–1869, 2007.
78. Fu W, Wei J, Gu J. MEF2C mediates the activation induced cell death (AICD) of macrophages. *Cell Res* 16:559–565, 2006.
79. Ralph P, Nakoinz I. Antibody-dependent killing of erythrocyte and tumor targets by macrophage-related cell lines: enhancement by PPD and LPS. *J Immunol* 119:950–954, 1977.
80. Stuehr DJ, Marletta MA. Synthesis of nitrite and nitrate in murine macrophage cell lines. *Cancer Res* 47:5590–5594, 1987.
81. Taniyama T, Holden HT. Cytolytic activity against tumor cells by macrophage cell lines and augmentation by macrophage stimulants. *Int J Cancer* 26:61–69, 1980.
82. Carvalho Neta AV, Stynen AP, Paixao TA, Miranda KL, Silva FL, Roux CM, Tsois RM, Everts RE, Lewin HA, Adams LG, Carvalho AF, Lage AP, Santos RL. Modulation of the bovine trophoblastic innate immune response by *Brucella abortus*. *Infect Immun* 76:1897–1907, 2008.